

SH3 Domain–Based Phototrapping in Living Cells Reveals Rho Family GAP Signaling Complexes

Hirokazu Okada,¹ Akiyoshi Uezu,¹ Frank M. Mason,¹ Erik J. Soderblom,² M. Arthur Moseley III,² Scott H. Soderling^{1*}

Rho family GAPs [guanosine triphosphatase (GTPase) activating proteins] negatively regulate Rho family GTPase activity and therefore modulate signaling events that control cytoskeletal dynamics. The spatial distribution of these GAPs and their specificity toward individual GTPases are controlled by their interactions with various proteins within signaling complexes. These interactions are likely mediated through the Src homology 3 (SH3) domain, which is abundant in the Rho family GAP proteome and exhibits a micromolar binding affinity, enabling the Rho family GAPs to participate in transient interactions with multiple binding partners. To capture these elusive GAP signaling complexes in situ, we developed a domain-based proteomics approach, starting with in vivo phototrapping of SH3 domain-binding proteins and the mass spectrometry identification of associated proteins for nine representative Rho family GAPs. After the selection of candidate binding proteins by cluster analysis, we performed peptide array–based high-throughput in vitro binding assays to confirm the direct interactions and map the SH3 domain-binding sequences. We thereby identified 54 SH3-mediated binding interactions (including 51 previously unidentified ones) for nine Rho family GAPs. We constructed Rho family GAP interactomes that provided insight into the functions of these GAPs. We further characterized one of the predicted functions for the Rac-specific GAP WRP and identified a role for WRP in mediating clustering of the postsynaptic scaffolding protein gephyrin and the GABA_A (γ-aminobutyric acid type A) receptor at inhibitory synapses.

INTRODUCTION

The Rho family small guanosine triphosphatases (GTPases), which include Rho, Rac, and Cdc42, function as molecular switches that cycle between guanosine triphosphate (GTP)–bound, active forms and guanosine diphosphate (GDP)–bound, inactive forms. In the active state, Rho family GTPases bind to effectors, such as kinases, formins, and the family of Wiskott-Aldrich syndrome proteins (WASPs), to form distinct actin cytoskeletal structures that include linear bundled and branched networks of filaments. By governing the assembly and disassembly of these actin structures, Rho family GTPases regulate various cellular activities such as cell migration, cell differentiation, cytokinesis, intracellular membrane trafficking, angiogenesis, and neuronal morphogenesis such as axonal guidance and synapse formation (1, 2). GEFs (guanine nucleotide exchange factors) promote the activation of Rho family GTPases by promoting the exchange of GDP for GTP, whereas GAPs (GTPase-activating proteins) turn them “off” by stimulating the intrinsic GTP-hydrolyzing activity of the GTPases to accelerate their conversion to the inactive form. Characterization of individual GAPs indicates that their activities are regulated by various mechanisms including protein-protein interaction, lipid binding, subcellular translocation, phosphorylation, and proteolytic degradation (3). GAPs for Rho family GTPases (Rho family GAPs) are composed of multiple modular domains, most of which have been identified as protein-protein interaction domains, lipid-binding domains, or enzymatic domains (4). This suggests that GAP interactions with proteins or lipids may mod-

ify Rho family GAP activities in coordination with other signaling pathways. SH3 (Src homology 3) domains are the most prevalent protein-protein interaction domains found in Rho family GAPs. Despite the likely importance of Rho family GAP SH3 domain interactions on Rho GTPase signaling, their binding partners are mostly unknown.

SH3 domains have a micromolar-range binding affinity [dissociation constant (K_d) of 1 to 200 μ M] (5, 6). SH3 domain interactions with ligand can be transient and depend on the cellular context—such as stimuli that affect ligand availability (6, 7). Because SH3 domains have a relatively broad ligand selectivity motif, each SH3 domain is likely to interact with several ligand proteins in vivo. These competitive interactions are thought to enable crosstalk between signaling pathways and result in a high degree of connectivity within a pathway. Such weak and transient interactions may therefore be crucial to the cellular roles of SH3 domain-containing GAP proteins. Conventional affinity purification methods may fail to detect transient or weak interactions because of cell extract preparation procedures or extensive washing. We performed in vivo phototrapping of SH3 domain-interacting proteins with a photoreactive amino acid cross-linker genetically incorporated into the proximity of the ligand-binding pocket of SH3 domains to identify SH3-ligand interactions in situ. We purified the cross-linked SH3-ligand complexes by a tandem affinity purification (TAP) strategy and identified the ligands by mass spectrometric analyses. We applied this methodology to the SH3 domains of 9 of the 14 human SH3 domain-containing Rho family GAPs. The interactions were verified by peptide array–based in vitro binding assays and coimmunoprecipitation experiments. Finally, we constructed protein interaction maps for each GAP to obtain mechanistic insights into their possible cellular functions. We found that WRP [WAVE (WASP family verprolin-homologous protein)–associated Rac GAP] binds to the principal inhibitory synapse scaffolding protein, gephyrin, and facilitates the postsynaptic clustering of gephyrin

¹Department of Cell Biology, Duke University Medical School, Durham, NC 27710, USA. ²Institute for Genome Science & Policy Proteomics Core Facility, Duke University Medical School, Durham, NC 27710, USA.

*To whom correspondence should be addressed. E-mail: s.soderling@cellbio.duke.edu

and ionotropic γ -aminobutyric acid type A (GABA_A) receptors. These results demonstrate the physiologic relevance of interactions identified by our approach and suggest that the WRP and gephyrin interaction may provide a mechanism to facilitate organization of receptors and signaling proteins at inhibitory synapses.

RESULTS

Photo-cross-linking of ligands for Rho family GAP SH3 domains

To identify Rho family GAP-containing protein complexes organized by SH3 domains, we developed a multistep workflow (Fig. 1A) based on an initial screen that used photo-induced trapping of intracellular interactions. *p*-Benzoyl-L-phenylalanine (pBpa) is a phenylalanine derivative that is a highly efficient and highly specific photo-activatable cross-linker (8, 9) that can be translationally inserted into proteins of interest in mammalian cells (10–14). pBpa can be incorporated into proteins of interest in a site-specific manner through the concomitant expression of the pBpa-specific

variant of *Escherichia coli* tyrosyl-tRNA^{Tyr} (*Eco*-pBpaRS) and *Bacillus stearothermophilus* suppressor tRNA^{Tyr} (Fig. 1B). We cloned Rho family GAP SH3 domains into a mammalian expression vector that coexpressed *Eco*-pBpaRS and the suppressor tRNA^{Tyr} and covalently trapped protein interactions in cells transfected with this construct by flowing suspended cells through a chamber with a 350- to 365-nm light source (Fig. 1C). We attached a TAP tag to the SH3 domain to enable the purification of the resulting SH3 domain–ligand complexes with a high signal-to-noise ratio (Fig. 1D). We incorporated pBpa into nine SH3 domains (Fig. 2A). To effectively cross-link the SH3 domain to ligands, pBpa must be located near the ligand-binding pocket; however, pBpa should not replace an amino acid directly involved in the binding interaction. We selected two candidate amino acid positions on the basis of structural data for the SRGAP1 SH3 domain (Fig. 2B) (15). These sites were proximal to the ligand-binding pocket but showed little conservation in sequence alignments, suggesting that they are less likely to be critical for ligand interactions (Fig. 2A, yellow highlights). To investigate whether pBpa successfully cross-linked to potential SH3 ligands in vivo, we expressed the modified SRGAP2 SH3 domain in mammalian cells in the presence of pBpa. Exposure to ultraviolet




Fig. 1. Experimental design to identify Rho family GAP protein complexes. (A) Workflow of overall approach to construct Rho family GAP interactomes. (B) Schematics for in vivo phototrapping strategy. A photo-activatable cross-linker, pBpa, is translationally incorporated into a cellular protein of interest at the site designated by an amber codon when co-expressed with a pBpa-specific tRNA synthetase and amber suppression tRNA. pBpa covalently cross-links with a binding protein when UV light is applied. (C) Modified cross-linking chamber for cells in suspension with inlet, outlet, and 9-W 350- to 365-nm light

source. (D) Schematic of the SH3 domain expression construct. TAP tag (protein A, TEV protease cleavage site, and CBP) was attached to the SH3 domain for the effective isolation of SH3 domain–ligand cross-linked protein complexes. A V5 epitope was added to follow the TAP purification process.

(UV) light produced a number of high-molecular weight species, indicating that light induced pBpa-mediated cross-linking of ligands to the SH3 domain (Fig. 2C), whereas no cross-linking was apparent in the absence of light (Fig. 2D). To purify the cross-linked complexes, we transfected FreeStyle 293 cells with the modified SRGAP2 SH3 domain and added pBpa (1 mM). Three days after transfection, we exposed the cells to 350- to 365-nm light in a modified cross-linking chamber to induce phototrapping of potential SH3 ligands, lysed them, and isolated the SH3 domain and co-purifying proteins by TAP (Fig. 2E). We subjected the purified samples to tryptic digestion and identified the proteins linked to the SH3 domain by liquid chromatography and tandem mass spectrometry (LC-MS/MS).

Hierarchical clustering of MS-identified SH3-interacting proteins

Expression of the nine SH3 domains was rescued by coexpression of the suppressor transfer RNA (tRNA) and tRNA synthetase, although the abundance of the different proteins varied (fig. S1). SH3 expression, cross-linking, and purification were performed in duplicate with the nine Rho family GAP SH3 domains shown in Fig. 2A, and interacting proteins were identified by LC-MS/MS analysis. A total of 860 interacting proteins were identified with a stringency scoring threshold [1.6% peptide and 14.2% protein false discovery rates (FDRs)] aimed at minimizing the number of potential false negatives likely to occur at higher stringency. This lower-stringency candidate interactor list was used as the basis for




Fig. 2. In vivo phototrapping and purification of ligands for Rho family GAP SH3 domains. (A) Sequence alignment of the Rho family GAP SH3 domains used in this study. Two amino acid positions (in yellow) were used to place pBpa within the SH3 domains listed. β Sheets a to e of the SH3 domain structure are indicated above. (B) Positions of pBpa within the structure of a Rho family GAP SH3 domain. The SH3 domain structure of SRGAP1 was used to determine the candidate residues (in yellow) to be replaced with pBpa that correspond to the residues in yellow in (A). Ligand-binding pocket is shown in blue. (C) In situ photo-cross-linking of an SH3 domain. Cells expressing V5-tagged wild-type (WT) SRGAP2 SH3 domain (lanes 1 and 4) or amber mutations (D764amb: lanes 2 and 3; R745amb: lanes 5 and 6) were grown with (lanes 2 and 5) or without (lanes 3 and 6)

pBpa and subjected to UV light. Western blot analysis of the immunoprecipitates by anti-V5 antibody reveals cross-linked high-molecular weight SH3-linked protein complexes. Representative blot from $n = 2$ is shown. (D) Photoactivation-dependent cross-linking of an SH3 domain. Cells expressing TAP-tagged SRGAP2 R745amb mutation SH3 domain were incubated with pBpa. After 3 days, cells were exposed to UV light in a photo-cross-linking chamber. Western blot analysis reveals photo-activated cross-linking of the SH3 domain. Representative blot from $n = 2$ is shown. (E) TAP of cross-linked SH3 domain protein complexes. The cell lysate obtained in (D) was subjected to TAP. The final eluate was concentrated by centrifugal filtration (Conc. sample) and subjected to mass spectrometric analysis. Representative blot from $n = 3$ is shown.

additional interaction validations to generate higher-confidence interaction data (see below). The relative abundance of the candidate SH3 domain interactors in each sample was evaluated by a modified spectral counting method (16, 17). To decode the pattern of associated proteins for each of the nine bait SH3 domains, we hierarchically clustered the identified proteins with unbiased Pearson correlation based on the mean normalized spectral counts (Fig. 3A) (18). This clustering enabled a rapid categorization of the proteins into two groups: one group that was predominantly associated with a single SH3 domain and a second group that showed no propensities toward any specific bait. The second type likely represents nonspecific or false positives that may be common contaminants with all baits. We therefore focused on the first type of clusters, which we called protein interaction (PI) clusters, for further analyses (Fig. 3B). The association between each SH3 domain bait and the proteins in the corresponding

PI cluster is of relatively high confidence because the other eight SH3 domain baits serve as negative controls. From the initial pool of 860 candidate proteins, we assigned 563 to PI clusters, substantially reducing the number of proteins to be further characterized. We assessed the abundance of proteins from the input cell lysate over a range of about three orders of magnitude (fig. S1C). Of the 563 proteins assigned to the nine PI clusters, only 10.8% were found in the input lysate (fig. S1D). Thus, most of the proteins identified in PI clusters appear to be low-abundance proteins that were enriched during the phototrap-based purification procedure.

SH3 domain ligand mapping by peptide array-based in vitro binding assay

The PI clusters contained not only proteins directly cross-linked to SH3 domains but also proteins that were stably associated with these cross-linked




Fig. 3. Elucidation of protein clusters enriched with Rho family GAP SH3 domains after in vivo phototrapping. (A) Hierarchical clustering of MS-identified SH3 domain-associated proteins. Hierarchical clustering analysis was performed with unbiased Pearson correlation of the mean normalized spectral counts for the Rho family GAP SH3 domains listed in Fig. 2. Protein clusters designated with blue bars contain proteins that predominantly

associated with individual SH3 domains. The degree of correlation in each cluster is shown. (B) Protein cluster with a predominant specificity for the ARHGAP26 SH3 domain. Proteins that have relatively high spectral counts or have been implicated in cytoskeletal regulation are exhibited. This protein cluster contains two proteins (PTK2 and PKN3 in orange) known to interact with ARHGAP26 in an SH3 domain-dependent manner.

proteins. To identify those proteins that bound directly to the SH3 domains and to validate these interactions, we performed high-throughput *in vitro* binding assays (Fig. 4 and fig. S2, A to I). SH3 domains bind to proline-rich sequences containing a PXXP motif (where P is Pro and X is any amino acid) (6, 19). Because Rho family GTPases play a prominent role in regulating the cytoskeleton, we selected candidate proteins from the PI clusters that either had been implicated in cytoskeletal regulation or showed high

spectral counts relative to other proteins in the cluster, and searched the sequences of those proteins for PXXP motifs. Peptides (18-mer) containing the PXXP motifs were synthesized on polyethylene glycol (PEG)-based cellulose membrane, incubated with purified glutathione *S*-transferase (GST)-tagged SH3 domains, and immunoblotted with an anti-GST antibody, and binding strength, normalized to the strongest interaction for each SH3 domain, was quantified (fig. S2, A to I). Peptides that bound

Fig. 4. Identification of SH3 domain ligands. (A) Array-based binding assay to identify SH3 domain-binding peptides. Peptides (18-mer) containing PXXP motifs from proteins in the ARHGAP26 PI cluster were synthesized, incubated with purified GST-tagged ARHGAP26 SH3 domain, and immunoblotted with anti-GST antibody. Colored rectangles and numbers indicate the positions of peptides that correspond to the peptide sequences listed in (B). Representative blot from $n = 2$ is shown. (B) SH3 ligand identification and determination of relative binding strength. Peptide immunoblots generated in (A) were densitometrically quantified. The apparent binding strength was normalized to the strongest interaction for ARHGAP26. PXXP motifs are indicated in orange. Numbers indicate unique motifs bound to the ARHGAP26 SH3 domain with more than 5% normalized binding strength. Note that peptides overlap each other (indicated in blue) and contain the identical core motif sequences. See fig. S2, A to I, for the *in vitro* binding assays for all Rho family GAP SH3 domains analyzed. (C) Coimmunoprecipitation (IP) of full-length ARHGAP26 by MICAL1. The cell lysates of HEK293 cells expressing GFP-tagged ARHGAP26 with or without V5-tagged MICAL1 were immunoprecipitated and blotted as indicated. Representative blot from $n = 2$ is shown. (D to F) SH3 domain ligands and their binding strengths for SRGAP2, ARHGAP4, and SNX26. PI clusters specific to SRGAP2, ARHGAP4, or SNX26 were subjected to peptide array-based *in vitro* binding assays as for ARHGAP26. Peptides that showed positive binding (more than 5% normalized binding strength) to each SH3 domain are shown. (G to I) Coimmunoprecipitation experiments using full-length Rho family GAPs. (G) HEK293 cells coexpressing gephyrin-V5 with vector or SRGAP2-Flag. (H) GFP-WAS2 with vector or ARHGAP4-Flag. or (I) SNX26-Myc with CDC42BPA-Flag or vector were subjected to immunoprecipitation and Western blot analysis by indicated antibodies. $n = 2$ to 4 for each coimmunoprecipitation. See fig. S2, J and K, for additional coimmunoprecipitation experiments.




Table 1. Direct interacting proteins for each SH3 domain. Peptide sequences that bound to each SH3 domain (based on the peptide array interaction screens). Protein names and corresponding gene nomenclature as well as relative binding strengths are listed. PXXP motif for each peptide is shown in orange.

Protein name	Gene name	Accession	SH3 binding sequence	Relative binding strength (%)	Cytoskeletal function
ARHGAP26					
Serine-threonine protein kinase N3	<i>PKN3</i>	Q6P5Z2	EMTP PPKPP RLYLP QE PT	100	Focal adhesion turnover Actin depolymerization
Focal adhesion kinase 1	<i>PTK2</i>	Q05397	D PA APP KKPPR PGAPGHL	91	
NEDD9-interacting protein with calponin homology and LIM domains	<i>MICAL1</i>	Q8TDZ2	DPEME PPKPP RSCSALA	64	
Ankyrin repeat and SAM domain-containing protein 1A	<i>ANKS1A</i>	Q92625	DRPYEEP PQKPP RFSQLR	27	
Rab11 family interacting protein 5	<i>RAB11FIP5</i>	Q9BXF6	RIMETSPTLLQI PPGPPK	6	
ARHGAP10					
Paxillin-associated protein with ARF GAP activity 3	<i>ASAP2</i>	O43150	GISQIR PPLPPQPP SRL	100	Arf-GAP. Recruits paxillin to focal contacts Dynein heavy chain Focal adhesion component
Dynein heavy chain 6	<i>DNAH6</i>	Q9C0G6	EKER PPKPEAP WLPTATW	95	
Talin-1	<i>TLN1</i>	Q9Y490	RIIRERI PEAPAG PPSDF	80	
Large proline-rich protein BAT3	<i>BAT3</i>	P46379	QTLGQQV PGFPTA TRVV	58	
Protein Shroom 3	<i>SHROOM3</i>	Q8TF72	ASDSGRG PQRP DARLLRS	37	
Protein phosphatase 1B	<i>PPM1B</i>	O75688	FSTQDHK PCNP REKERIQ	19	F-actin binding. Apical constriction
ARHGAP10L					
Myosin-14	<i>MYH14</i>	Q7Z406	AQPFLFT PRGPS AGGGPG	100	Myosin heavy chain F-actin binding Interacts with the formin-regulatory protein Flightless
Drebrin	<i>DBN1</i>	Q16643	AAIIAQR PDNP REFFKQQ	68	
Leucine-rich repeat flightless-interacting protein 2	<i>LRRFIP2</i>	Q9Y608	YNGGLYN PYGPRT PSECS	65	
FH1/FH2 domain-containing protein 3	<i>FHOD3</i>	Q2V2M9	HREA PGPPPPPP TFLGL	53	
LIM domain only protein 7	<i>LMO7</i>	Q8WWI1	LYVRKLS PVMPN PGNAFD	12	
LIM homology domain- and calponin homology domain-containing protein 1	<i>LIMCH1</i>	Q9UPQ0	G PFSPCSPTPP GQSPNRS	11	Contains actin binding calponin homology domain
ARHGAP12					
U1 small nuclear ribonucleoprotein C	<i>SNRPC</i>	P09234	GAMIP PPPSLPGPPR PGM	100	Tight junction protein Controls junctional ion transport by inhibiting WNK4
Tight junction protein ZO-2	<i>TJP2</i>	Q9UDY2	FGRSILK PSTPIP QEGE	93	
Serine-threonine protein kinase WNK1	<i>WNK1</i>	Q9H4A3	NATALEL PGLPL SLPQPS	52	
Ubiquitin-associated protein 2-like	<i>UBAP2L</i>	Q14157	YSIPFPT PTT PLTGRDGS	45	
Phosphoribosylformylglycinamide synthase	<i>PFAS</i>	O15067	FPKASV PREPGG SPRVA	24	
ARHGAP4					
Diaphanous homolog 1	<i>DIAPH1</i>	O60610	LPGGTAI PPPPPL GSAR	100	Formin. Actin nucleation and elongation
Splicing factor 3B subunit 2	<i>SF3B2</i>	Q13435	LQ PPPPPPPPPP GLGLGF	68	
Radixin	<i>RDX</i>	P35241	LKTVMSA PPPPPPPPVIP	53	Cross-link F-actin to plasma membrane
Trifunctional enzyme subunit β	<i>HADHB</i>	P55084	FRFNFLA PELPA VSEFST	44	
Wiskott-Aldrich syndrome protein family member 2	<i>WASF2</i>	Q9Y6W5	PGFAP PPAPPPPPPP MIG	41	Arp2/3-mediated actin nucleation promoter
DAZ-associated protein 1	<i>DAZAP1</i>	Q96EP5	QFSFGYG PPPPPPD QFAP	18	
Splicing factor 1	<i>SF1</i>	Q15637	GMPPFGM PPAPPPPP QNN	7	
SRGAP2					
YLP motif-containing protein 1	<i>YLPM1</i>	P49750	HHL PPPLPPPPVMP GGG	100	Inhibitory postsynaptic scaffold protein. Associates with cytoskeleton
Gephyrin	<i>GPHN</i>	Q9NQX3	FGRVFMK PGLP TTTFATLD	55	

continued on next page

Protein name	Gene name	Accession	SH3 binding sequence	Relative binding strength (%)	Cytoskeletal function
Palladin	<i>PALLD</i>	Q8WX93	FPLPPPPPLPSPGQASH	43	Actin remodeling and bundling
Glutamate–cysteine ligase catalytic subunit	<i>GCLC</i>	P48506	WQTMRFKPPPPNSDIGWR	18	
Dishevelled, dsh homolog 3	<i>DVL3</i>	Q92997	MAFPRPRPWGPQEPLRAA	9	
ARHGAP9					
SET and MYND domain–containing protein 3	<i>SMYD3</i>	Q9H7B4	NGLRAVTPLRPGELLFRS	100	Complexes with BAIAP2 and regulates actin dynamics
Coiled-coil domain–containing protein KIAA1826	<i>KIAA1826</i>	Q8NCY6	SRRENELPDFPHIDEFFT	70	
EGFR kinase substrate 8	<i>EPS8</i>	Q12929	PKEQFIPPYVPRFRNGWE	24	
UPF0027 protein C22orf28	<i>C22orf28</i>	Q9Y3I0	GSTRAFPHHPLIAVDYQ	13	
SNX26					
Proline-rich protein 4	<i>PRR4</i>	Q16378	GHRQLSLPRFPVSVLQEA	100	Cdc42-activated serine-threonine kinase
Serine-threonine protein kinase MRCK α	<i>CDC42BPA</i>	Q5VT25	IQILKDLPMNRPQESRT	60	
Statherin	<i>STATH</i>	P02808	RFGYGYGPYQVPPEQPLY	28	Cytoskeletal remodeling Actin binding. Regulates filament dynamics
Tyrosine protein kinase ABL1	<i>ABL1</i>	P00519	KKKKKTAPTPPKRSSSFR	20	
Adenylyl cyclase–associated protein 1	<i>CAP1</i>	Q01518	PPPPPPCPPPPPVSTISC	10	
RICS					
Cathepsin B	<i>CTSB</i>	P07858	CGTFLGGPKPPQRMVTE	100	Proteolysis of substrates involved in cytoskeletal remodeling
Chromodomain helicase-DNA binding protein 3	<i>CHD3</i>	Q12873	YLNLSQEPAAHPAMALHAR	88	
Protein FAM134C	<i>FAM134C</i>	Q86VR2	ALVEVLGPYEPLLSRVQA	85	
Calpain-2 catalytic subunit	<i>CAPN2</i>	P17655	AGTLFQDPSFPAIPSALG	49	
Serotransferrin	<i>TF</i>	P02787	NLREGTCPEAPTDECKPV	49	
Inositol-3-phosphate synthase 1	<i>ISYNA1</i>	Q9NPH2	LSFLFKAPLVPPGSPVVN	45	
Crossover junction endonuclease MUS81	<i>MUS81</i>	Q96NY9	RSRPWGTGNPESGAMTS	31	
Eukaryotic translation initiation factor 2- α kinase 3	<i>EIF2AK3</i>	Q9NZJ5	VRISEKFPSSPKALESVT	15	
CDK5 and ABL1 enzyme substrate 1	<i>CABLES1</i>	Q8TDN4	GCIALAAPGTPAAGLAAG	7	Facilitates CDK5 activation
SRA stem-loop–interacting RNA binding protein, mitochondrial	<i>SLIRP</i>	Q9GZT3	VQVHTRRPKLPQTSDDK	7	
Vigilin	<i>HDLBP</i>	Q00341	PTYKDAFPPLPEKAACLE	5	

within the relative range of 5 to 100% of the strongest interaction for each SH3 domain were scored as positive. In the case of ARHGAP26, we synthesized 91 unique peptides from the 18 proteins listed in Fig. 3B in an array format [3 of the 18 proteins, C3orf10 (probable protein brick1), CYFIP1 (cytoplasmic FMR1-interacting protein 1), and NCKAP1 (Nck-associated protein 1), had no PXXP motifs]. The SH3 domain overlay showed binding to only nine unique peptides (Fig. 4, A and B), indicating that the SH3 domain interaction is highly specific. These peptides corresponded to sequences from five different proteins, MICAL1 (molecule interacting with CasL 1), PTK2 [protein tyrosine kinase 2, also known as focal adhesion kinase (FAK1)], ANKS1A (ankyrin repeat and sterile α motif domain–containing 1A), PKN3 (protein kinase N3), and RAB11FIP5 (Rab11 family interacting protein 5). The SH3 domain of ARHGAP26 is known to bind PTK2 and PKN3 (20, 21). Thus, the phototrapping of bound ligands with pBpa-modified SH3 domains retains the ligand preference of the wild-type domains and, moreover, can recapitulate known interactions. We further validated the interaction between ARHGAP26 and MICAL1, which, on the basis of spectral counts, was the most enriched protein in the ARHGAP26 PI cluster, in a co-immunoprecipitation assay with full-length constructs (Fig. 4C).

Analysis of peptides that bound directly to other GAP SH3 domains revealed multiple previously unidentified interactions (Fig. 4, D to F, and fig. S2, A to I). To verify that full-length proteins could recapitulate the SH3-peptide interactions, we tested several interactions by coimmunoprecipitation using full-length constructs. We confirmed the peptide-based identification of interactions between Rho family GAPs and multiple types of proteins: (i) scaffold proteins gephyrin and SRGAP2 (Fig. 4G), palladin and SRGAP2 (fig. S2J); (ii) activators of actin polymerization WASF2 (Wiskott-Aldrich syndrome family protein 2, also referred to as WAVE2) and ARHGAP4 (Fig. 4H), DIAPH1 (diaphanous homolog 1) and ARHGAP4 (fig. S2K); and (iii) kinases activated by members of the Rho family, CDC42BPA (myotonic dystrophy kinase–related Cdc42-binding kinase) and SNX26 (Fig. 4I). These data confirmed that domain-based interactions can reproducibly identify complexes that exist between full-length proteins (22). Overall, 789 unique peptides from 163 proteins were screened with the peptide array. Positive binding, indicating direct ligands for the Rho family GAPs, was detected for 102 peptides (13%) from 54 proteins (33%). We identified 4 to 7 direct protein interactions for each SH3 domain except for RICS [Rho family GAP involved in β -catenin–N-cadherin

and NMDA (*N*-methyl-D-aspartate) receptor signaling, also known as p250GAP], for which we identified 11 direct protein interactions. The result of the peptide array analyses, showing which candidate proteins interacted directly with each SH3 domain and providing the SH3 binding sequence for each protein, is summarized in Table 1.

Bioinformatic analyses to construct Rho family GAP interactomes

We next examined whether the proteins that interacted directly with each SH3 domain were likely to interact with other proteins in the corresponding PI clusters. Bioinformatic analysis was used to combine the direct SH3 domain interactions identified here (brown or orange edges) with previously known interactions (blue edges) among the proteins in the PI clusters (represented by circle nodes) (see Materials and Methods) (Fig. 5A and fig. S3, A to I). As illustrated for ARHGAP26 [Fig. 5A (left)], many of the proteins in the PI cluster interacted with one another, although most were only indirectly associated with the pertinent Rho family GAP. To identify possible cellular functions for each Rho family GAP, we characterized these interactomes further by focusing on the direct or primary binding interactions with each Rho family GAP and those proteins that were known to interact with these primary binders [Fig. 5, A (right) and B to I]. Cellular functions of some of proteins that interact directly with the SH3 domain (see Table 1) and their associated proteins have been identified, allowing us to infer the cellular functions in which each GAP may be involved. In most cases, multiple proteins that bound directly to a particular GAP had been implicated in the same functions, reinforcing the functional implications. Highlights from this network analysis are summarized below.

The SH3 domains of ARHGAP26 and ARHGAP10 are 68% identical and directly bind PTK2, a key kinase that promotes the turnover of focal adhesions, which are protein complexes that link extracellular matrix binding to signaling to the cytoskeleton (20, 23). We reproducibly identified ARHGAP26 (Fig. 5A, right) as a PTK2-binding protein in the present study. ARHGAP10 (Fig. 5B) instead bound to distinct focal adhesion elements: talin (TLN1), a critical structural component of the focal adhesion complex and ASAP2 [ArfGAP with SH3 domain, ankyrin repeat, and PH (pleckstrin homology) domain 2], which is a negative regulator of PXN (paxillin, an adaptor protein that organizes PTK2 and other proteins at focal adhesions) recruitment to focal contacts (24). These results suggest that ARHGAP26 and ARHGAP10 play different roles in focal adhesion assembly. ARHGAP26 also bound tightly to MICAL1, an actin filament disassembly factor activated downstream of semaphorin-plexin signaling (25), suggesting that ARHGAP26 may play a role in axonal pathfinding.

ARHGAP10L (Fig. 5C) bound to six proteins, including FHOD3 (a myocardial formin; formins stimulate actin polymerization downstream of Rho family GTPases), LRRFIP2 [leucine-rich repeat (in FLII) interacting protein], LMO7 (LIM domain only protein 7), and MYH14 (myosin14), all of which are highly expressed in skeletal muscle and heart. LRRFIP2 is associated with Flightless-1 (Fli-1), which enhances the activity of formin family proteins (26), implying a role of ARHGAP10L in FHOD3-mediated actin assembly that may be stimulated by Fli-1. LMO7 is required for normal heart development in zebrafish (27).

ARHGAP12 (Fig. 5D) localizes to cell-cell junctions in mouse epithelial tissues, including small intestine, kidney, salivary gland, and liver (28), but its function remains unknown. We found that ARHGAP12 bound to the tight junction component TJP2 [tight junction protein 2 or zona occludens 2 (ZO2)]. Immunostaining of ARHGAP12 in human bronchial epithelial cells showed colocalization with TJP2 at tight junctions (fig. S3J). ARHGAP12 also bound to the kinase WNK1 (lysine-deficient kinase 1), which inhibits WNK4, a negative regulator of ion transport through tight

junctions (29). These data suggest that ARHGAP12 may participate in tight junction formation and junctional ion transport.

ARHGAP4 (Fig. 5E), which is predominantly expressed in hematopoietic cells (30), bound to two distinct types of actin assembly activators, WASF2 and DIAPH1 (see Fig. 4H and fig. S2K for full-length coimmunoprecipitation). Moreover, ARHGAP4 SH3 domain phototrapping identified most members of WASF2 complex [CYFIP1, cytoplasmic fragile X interacting protein 1), NCKAP1, ABI1 (Abl interacting protein 1), ACTR2 (actin-related protein 2), and ARPC2 (actin-related protein 2/3 complex subunit 2)], together with WASF2, as associated with ARHGAP4. DIAPH1 and WASF2 are also highly expressed in hematopoietic cells, where they are required for polarization and cell migration (31). These interactions indicate that ARHGAP4 negatively regulates actin polymerization in cells by forming complexes with WASF2 and DIAPH1, both of which are activated downstream of Rho family GTPases (31, 32). Consistent with this conclusion, the ARHGAP4-related GAPs, WRP and SRGAP2, bind to WASF1 (also commonly known as WAVE1) and the formin FMNL1 (formin-like 1 protein), respectively, and oppose their actin polymerization and severing functions (33, 34).

SRGAP2 (Fig. 5F), a neuronal GAP, showed strong binding to an inhibitory synapse postsynaptic scaffolding protein, gephyrin, and an actin filament-bundling and remodeling protein, palladin (see Fig. 4G and fig. S2J for full-length coimmunoprecipitation). See below for our investigation of the functional link between gephyrin and SRGAP2 or the homologous WRP.


ARHGAP9 (Fig. 5G) is predominantly expressed in peripheral blood leukocytes, spleen, and thymus (35); its WW domain binds to mitogen-activated protein kinases (MAPKs) competitively within the MAPK kinase (MAPKK) binding site (36). Cell-based assays have shown that the ARHGAP9 and MAPK interaction suppresses epidermal growth factor receptor (EGFR)-induced actin reorganization, presumably by blocking MAPK activation (36). Our analysis identified the EGFR-associated protein EPS8 (EGFR pathway substrate 8) as bound to ARHGAP9. In addition to its own barbed-end actin capping activity, EPS8 stimulates mitogenic signaling to actin remodeling through SOS-1 (son of sevenless homolog 1), which functions as a GEF for Ras and Rac (37). EPS8 also interacts with BAIAP2 (brain-specific angiogenesis inhibitor 1-associated protein 2, also known as IRSp53) and facilitates Cdc42-mediated actin bundling, thereby promoting filopodial protrusion (38). ARHGAP9 may therefore connect MAPK and EPS8 signaling through ARHGAP9 SH3 domain-based interactions with EPS8.

SNX26 (sorting nexin 26, also called TCGAP or NOMA-GAP) (Fig. 5H), which, in both immature and adult mice, is predominantly expressed in the brain, inhibits the Rho GTPases TC10 and Cdc42 (39). SNX26 binds the receptor NTRK1 (neurotrophic tyrosine kinase receptor 1, also referred to as TrkA) and suppresses nerve growth factor (NGF)-induced neurite outgrowth in a GAP-dependent manner (40, 41). NGF-induced NTRK1 activation causes ABL1 (c-Abelson kinase 1 or c-Abl)-mediated phosphorylation of the adaptor protein Crk at Tyr²²², resulting in the dissociation of paxillin and c-Abl from Crk (42), which is critical for NGF-induced neurite outgrowth (43). Our analysis of the SNX26 SH3 domain identified c-Abl as a direct interaction, indicating a link between NGF signaling and Rho GTPase-mediated regulation of the cytoskeleton. We also identified two proteins involved in cell motility as SNX26-binding proteins: CDC42BPA, which facilitates myosin-dependent cell motility (44), and CAP1 (adenylyl cyclase-associated protein 1), which promotes the recycling of cofilin and actin for rapid actin turnover (45). CAP1-dependent facilitation of rapid actin turnover drives cell motility and may also bind to the Cdc42 GEF MCF2 (MCF2 cell line transforming sequence) (45, 46). CAP1 is enriched in neuronal growth cones, and its depletion decreases growth cone F-actin content and impairs neuronal outgrowth (47, 48). In *Drosophila*, CAP1 genetically interacts with c-Abl during axonal guidance (49). Thus, SNX26 may act downstream of

Fig. 5. Rho family GAP interactome graphs with inferred cellular functions. Protein-protein interaction networks constructed from the interactions identified in this study and previously identified (physical) interactions. Circle nodes represent proteins in the PI clusters specific to individual SH3 domains. Hexagon nodes represent proteins that were associated with two SH3 domains but were excluded from PI clusters in the hierarchical clustering. Both nodes are colored in the purple spectrum (bottom left of each graph) reflecting the mean normalized spectral counts. Light blue octagon nodes indicate proteins identified in previous studies. Diamond nodes in cyan represent GTPases. Brown edges represent interactions identified here, and orange edges indicate known interactions reproduced in this study. Edge thickness reflects the relative binding affinity determined by *in vitro* binding assays. Blue and cyan edges of constant thickness indicate previously known interactions. (A, left) ARHGAP26 interactome graph. Proteins from the ARHGAP26-specific PI cluster (circle nodes) and ARHGAP26-enriched proteins (hexagon nodes) make a densely connected protein network. (See fig. S3, A to I, for other Rho family GAP interactomes.) (A to I) Graphs of Rho family GAP interactomes based on direct interactions identified in this study combined with previously known interactions to infer cellular functions. Subnetworks with specific cell functions are grouped by dashed line and indicated (see text for details).

NGF to coordinate TC10- and Cdc42-mediated responses with ABL1 and CAP1 signaling to promote neurite outgrowth.

RICS (Fig. 5I) is a brain-enriched GAP that inhibits dendritic spine formation and maturation (50, 51). RICS binds to subunit 2B (GRIN2B) of the NMDA-type glutamate receptor, and NMDA decreases RICS abundance. The NMDA-dependent decrease in RICS levels occurs principally through an increase in the microRNA miR132, which contributes to a decrease in RICS translation (52). We identified two proteases as RICS-binding proteins. One of these proteases, m-calpain (CAPN2), shows Ca^{2+} -dependent proteolytic activity upon NMDA-evoked Ca^{2+} influx, suggesting that RICS may also facilitate NMDA-dependent proteolysis of CAPN2 substrates within dendritic spines. Peptide array analysis also revealed that RICS bound to CABLES1 (Cdk5 and ABL1 enzyme substrate), a scaffolding protein for two kinases, CDK5 (cyclin-dependent protein kinase 5) and ABL1. NMDA stimulates CAPN2 to cleave the CDK5-associated cofactor p35 (CDK5R1), thereby activating CDK5 (53). Thus, RICS may facilitate CDK5 activation by promoting a complex between CAPN2, CABLES, and CDK5. RICS also bound to another protease CTSB



(cathepsin B), which may negatively regulate the abundance of another CABLES-associated protein, TP53 (tumor suppressor p53) (54).

Gephyrin clustering facilitated by WRP

To investigate the physiological relevance of the SH3 domain-mediated Rho family GAP interactions we identified, we characterized the interactions between SRGAP2 and the scaffolding protein gephyrin. Gephyrin contains two self-association domains, called the G and E domains (55). Together, these domains facilitate the organization of gephyrin clusters that anchor glycine and GABA_A receptors at postsynaptic sites along the dendritic shaft (55). The formation of gephyrin clusters appears to depend on interactions between the gephyrin E domain and GABA or glycine receptors and probably involves local regulation of the actin cytoskeleton (56, 57). We found that SRGAP2 bound to the gephyrin E domain, suggesting that SRGAP2 may promote gephyrin clustering *in vivo*.

To test this possibility, we first performed a coimmunoprecipitation analysis. We used an antibody directed against gephyrin to precipitate complexes from mouse brain extract (MBE) and immunoblotted with an antibody that recognizes both SRGAP2 and WRP, which share 90% identity in their SH3 domains (33, 34). SRGAP2 or WRP was readily detected in gephyrin immunoprecipitate from MBE, suggesting that SRGAP2, WRP, or both may interact with gephyrin *in vivo* (Fig. 6A). Coimmunoprecipitation of heterologously expressed WRP and gephyrin constructs from human embryonic kidney (HEK) 293 cells confirmed that WRP bound to gephyrin (fig. S4A); we also detected a weak association between gephyrin and WRP lacking the SH3 domain, indicating that indirect interactions between the two may also exist. *In vitro* peptide-binding assays revealed that the WRP and SRGAP2 SH3 domains bound to the same peptide in gephyrin (Fig. 6B). We had previously characterized WRP knockout mice as a potential model for 3p-syndrome mental retardation (a form of retardation resulting from microdeletions of chromosome 3p25-26) (58), and clustering gephyrin is induced by ARHGEF9, which is also associated with mental retardation (59, 60). Therefore, we focused on the potential role of WRP in gephyrin clustering. To assess WRP colocalization with gephyrin, we performed immunocytochemistry in HEK293 cells transfected with complementary DNAs (cDNAs) to express both. As previously shown in heterologous cells, gephyrin formed large clusters (Fig. 6C) (61). WRP colocalized with gephyrin at these clusters. Deletion of the WRP SH3 domain abolished colocalization, indicating that it depended on the SH3 domain (Fig. 6C). Because gephyrin clustering is relevant to its ability to cluster synaptic receptors, we examined the effect of WRP on the size of the gephyrin clusters. Coexpression of WRP with gephyrin increased the average size of gephyrin clusters by more than twofold compared to gephyrin alone (Fig. 6E). Distribution analysis showed that WRP decreased the fraction of clusters smaller than 4 μm^2 in area and increased the fraction of clusters larger than 6 μm^2 in area (Fig. 6F), suggesting that WRP induces or stabilizes gephyrin clustering. Immunocytochemical analysis of endogenous gephyrin in hippocampal neuronal cultures revealed punctate staining, which was mostly colocalized with VIAAT (vesicular inhibitory amino acid transporter), a presynaptic marker for inhibitory synapses (fig. S4B). Coexpression of green fluorescent protein (GFP)-tagged WRP increased the number of gephyrin clusters within the dendrite compared to that of GFP controls. Deletion of the SH3 domain abolished this effect, indicating that SH3 domain-dependent binding of WRP with gephyrin increased the dendritic clustering of gephyrin (Fig. 6, G and H).

Role of WRP in postsynaptic clustering of gephyrin and GABA_A receptors *in vivo*

To confirm a role for WRP in mediating gephyrin clustering at synapses *in vivo*, we performed comparative histological analyses of gephyrin stain-

ing in the hippocampi of wild-type and WRP-knockout mice (58). Gephyrin is present in pyramidal neurons in the CA1 region of hippocampal formation (62) (Fig. 7A). Therefore, we analyzed gephyrin staining within the stratum radiatum, where inhibitory synapses are formed on the dendrites of pyramidal cells. Consistent with the *in vitro* data, there was a decrease in the number of gephyrin clusters in WRP knockout mice compared to that of wild-type littermates (Fig. 7, B and C). Consistent with a role for gephyrin in promoting GABA_A receptor clustering, the number of GABA_A receptor puncta was decreased in WRP knockout mice (Fig. 7, B and D). Together, these data suggest that loss of WRP results in a decrease in gephyrin clustering and consequentially a reduction in GABA_A receptor synaptic puncta. To determine whether the gephyrin clusters we analyzed were synaptic, we co-stained with VIAAT. Most of the gephyrin puncta colocalized with VIAAT immunopositive clusters, indicating that the gephyrin immunostaining was mainly synaptic (fig. S5). Because the density of VIAAT puncta was unaffected in WRP knockout mice, deletion of WRP does not appear to alter inhibitory innervation. This is consistent with studies showing that the deletion of other gephyrin-associated proteins such as ARHGEF9 or Neuroligin2 does not alter inhibitory presynaptic innervation (62, 63). To further analyze the effect of WRP on individual gephyrin and GABA_A receptor clusters, we quantified the size of individual puncta. This analysis revealed that the average area of puncta was reduced for both gephyrin and GABA_A receptors in WRP knockout mice (Fig. 7F). Together, these results, when combined with the overexpression data, suggest that WRP promotes clustering of postsynaptic GABA_A receptors at inhibitory synapses.

DISCUSSION

Our study has provided insight into how Rho family GAPs may organize the formation of complex signaling networks through SH3 domain interactions. The methodology described here is likely applicable to any proteomics-based approach that uses proteins where structural information is available to guide the placement of the pBpa. Indeed, a wide range of protein binding events may be identified by this approach, including protein-DNA interactions (64). We used an overexpression strategy, which undoubtedly increases the abundance of the bait proteins relative to that found *in vivo*. In the future, this technique could be refined by genetically modifying the locus for endogenous proteins such that the unnatural amino acid approach could be used in primary cells without overexpression of protein baits. However, our results demonstrate that the SH3 domain-based phototrapping approach can reveal new insights into physiologically relevant Rho family GAP signaling pathways.

Methodology

Affinity purification-MS/MS (AP-MS/MS) is commonly used to identify protein-protein interactions and is often a first approach toward the analysis of the cellular functions of uncharacterized proteins (65). AP-MS/MS depends on stable interactions that can survive the multiple purification steps necessary for sample preparation. Many protein interactions are transient in nature or involve proteins of low abundance with micromolar binding affinities. These interactions may not be optimal for traditional AP-MS/MS approaches. Moreover, it is difficult to know whether interactions identified during standard purification approaches formed during lysis or actually occur within the intact cell. The methodology used here, in which protein interactions are covalently trapped in the intact cell, provides one approach for overcoming these limitations. Indeed, SH3 domains have a moderate (typically micromolar) binding affinity that enables multiple ligand interactions, some of which are likely to be transient. Only one protein identified by phototrapping and confirmed by peptide array




Fig. 6. WRP enhances gephyrin clustering in HEK cells and hippocampal neurons. (A) Coimmunoprecipitation of SRGAP2 and WRP by gephyrin in the mouse brain. Mouse brain extract (MBE) was subjected to immunoprecipitation with anti-gephyrin antibody. Coprecipitation of SRGAP2 or WRP (indicated by arrows) was observed by Western blot using anti-SRGAP2/WRP antibody. Representative blot from *n* = 4 is shown. (B) SRGAP2 and WRP bind the same site in gephyrin. Gephyrin peptides (18-mer) containing PXXP motifs were synthesized, incubated with purified GST-tagged SH3 domains of SRGAP2 or WRP, and immunoblotted with anti-GST antibody. Representative blot from *n* = 2 is shown. (C) Colocalization of WRP with clustered gephyrin in HEK293 cells. GFP-tagged gephyrin was coexpressed in HEK293 cells with V5-tagged WRP or WRP lacking its SH3 domain. Immunofluorescence was visualized with anti-GFP (green) and anti-V5 antibodies (red), or 4',6-diamidino-2-phenylindole (DAPI) (blue). Arrows indicate gephyrin clusters. Scale bars, 10 μm. (D to F) Increased size of gephyrin clusters upon WRP coexpression. Area of gephyrin clusters was measured in cells cotransfected with GFP-gephyrin and vector or WRP-V5. Average intensity (D), area (E), and area distribution (F) of gephyrin clusters from three independent experiments are shown. Data are presented as means ± SEM. **P* < 0.0005 (two-tailed *t* test). (G) SH3-dependent increase in endogenous gephyrin clustering by WRP. Hippocampal neurons transfected at DIV9 with GFP, GFP-tagged WRP, or WRP lacking the SH3 domain were fixed at DIV12 and stained with anti-gephyrin antibody. Scale bars, 10 μm (left panels) or 5 μm (right panels). (H) Quantification of the gephyrin clusters in (G). GFP, *n* = 42 neurons from seven mice; GFP-WRP, *n* = 27 neurons from four mice; GFP-WRP ΔSH3, *n* = 14 neurons from three mice. Data are presented as means ± SEM. ***P* < 0.0005 (two-tailed *t* test). N.S., not significant.

analysis was also identified in the input lysate data (Talin-1 from the ARHGAP10 PI cluster) (Fig. 5B and fig. S1D). Thus, the interactions we identified through this approach appear to represent those of lower-abundance proteins that were enriched by phototrapping. Furthermore, the successful detection of interacting proteins relied on UV-dependent covalent cross-linking, because without cross-linking we only identified nonspecific interactions.

In addition to using pBpa to entrap and identify SH3-dependent interactions, we used several additional steps to distinguish specific from nonspecific interactions. This included comparing the normalized abundance of proteins identified by MS from each SH3 domain bait to those of related SH3 domains. Proteins that were specifically enriched for an SH3 domain were considered to be candidate interacting proteins. This analysis reduced the number of possible interactions and enabled the generation of PI clusters for each SH3 domain. Further analysis using the peptide array-based binding assay revealed which of these proteins were likely to bind directly to each SH3 domain as well as the relative binding strengths and binding sites. These assays were done using the unmodified SH3 domain, indicating that binding interactions were not a result of altered specificity of the pBpa-containing domains. We also performed coimmunoprecipitation experiments with full-length proteins to validate a sample of the interactions we identified. All the interactions tested by coimmunoprecipitation were confirmed, suggesting that interactions identified by isolated SH3 domains were reliable. The success of this approach fits well with previous work showing that domain-based interaction studies can reliably recapitulate protein interactions that occur between full-length proteins (22).

By merging our interaction results for each SH3 domain with published protein association data, we obtained a more complete view of the possible protein networks organized around each Rho family GAP. These networks allowed us to predict possible functions for these GAPs based on the biological roles of their




Fig. 7. WRP promotes postsynaptic clustering of gephyrin and GABA_A receptors in vivo. **(A)** Coronal section of mouse brain. Immunohistological analysis was performed on the CA1 region of the hippocampal formation (boxed region) where inhibitory synapses are made on the dendrites of pyramidal neurons. **(B)** Reduction of gephyrin cluster density and associated GABA_A receptors in WRP knockout (KO) mice. Stratum radiatum of CA1 region in the hippocampal formation from WRP^{+/+} or WRP^{-/-} mice was stained with anti-gephyrin and anti-GABA_A receptor γ subunit antibodies. **(C and D)** Quantification of the density of gephyrin and GABA_A receptor puncta. *n* = 3 mice. From each mouse, four brain slices were processed for immunostaining and two images per slice were obtained. Data are presented as means ± SEM. ****P* < 0.0005; **P* < 0.01 (two-tailed *t* test). **(E)** Comparison of the average fluorescence intensity of gephyrin and GABA_A receptor puncta. **(F)** Decreased sizes of gephyrin and GABA_A receptor puncta in WRP knockout mice. Data are presented as means ± SEM. ***P* < 0.005 (two-tailed *t* test).

associated proteins. In most of the resulting networks, multiple proteins that had been previously implicated in the same functions were found for each Rho family GAP, increasing the likelihood that the predicted functional inference was correct. We also detected proteins in some networks that appeared to be outliers in that the function of these proteins did not seem to fit well with that of Rho family GAPs. These include histone regulatory protein (SMYD3 in Fig. 5G), splicing factors (SF1 and SF3B2 in Fig. 5E), or a negative regulator of telomerase activity (YLP1 in Fig. 5F) (66). These interactions may represent false positives or interactions that are outside of our current understanding of the possible functions of these GAPs. One unexpected function predicted by our study that we confirmed, how-

ever, was the role of SRGAP2 (or the closely related WRP) in the postsynaptic architecture of inhibitory synapses.

Inhibitory synapse regulation by WRP

Relatively little is known about the postsynaptic structural proteins at inhibitory synapses. However, gephyrin, the most abundant scaffolding protein at inhibitory synapses, has long been thought to mediate the formation and maintenance of the postsynaptic organization of receptors and signaling proteins (55). Gephyrin contains three major domains (G, C, and E domains). The G and E domains are believed to associate to form stable trimers or dimers, facilitating the clustering of gephyrin at postsynaptic sites that is also facilitated by other factors such as ARHGEF9 (55). However, the mechanisms that modulate the formation of these gephyrin clusters are not fully understood. We found that the Rac-specific GAPs SRGAP2 and WRP interact with gephyrin in vitro and in vivo. This interaction is likely to be physiologically relevant, because WRP increased the size and number of gephyrin clusters when overexpressed in HEK293 cells or in dissociated hippocampal neurons, whereas cluster size and number were decreased when WRP was knocked out in vivo. Our peptide array mapping study indicated that the SH3 domain of WRP binds to the gephyrin E domain, one of the domains responsible for gephyrin clustering. Both the actin filament and microtubule cytoskeleton have been implicated in gephyrin clustering (56, 57), and several cytoskeletal regulatory proteins interact with gephyrin, including the Cdc42 ARHGEF9 (55). ARHGEF9, by binding with neuroligin 2, promotes the translocation of gephyrin to the membrane (60–62, 67). Mice lacking ARHGEF9 exhibit region-specific decreases in the number of GABAergic inhibitory synapses (63). However, neither ARHGEF9 GEF activity nor Cdc42 signaling plays a role in synaptic gephyrin clustering (68). Our results suggest that Rac

signaling may contribute to gephyrin clustering, however, because WRP is selective for this GTPase (34). WRP has been implicated in a human 3p-syndrome mental retardation that in some, but not all, cases is associated with seizures (69). Mutations in WRP are also associated with schizophrenia, aspects of which may also be linked to altered GABAergic synaptic transmission (70–72). Thus, our discovery that loss of WRP in mice affects the postsynaptic structure of inhibitory synapses suggests a possible mechanism that could contribute to these neurological disorders associated with WRP mutations.

In summary, the data outline a new multistep approach for domain-based proteomics within cells that is applicable to many types of protein

interactions. Using this approach, we identified possible roles for Rho family GAPs in numerous aspects of cell physiology, including the clustering of gephyrin at inhibitory synapses.

MATERIALS AND METHODS

Plasmids

A mammalian expression vector that coexpresses the pBpa-specific variant of *Eco*-pBpaRS and *B. stearothermophilus* suppressor tRNA^{Tyr} (a gift of S. Yokoyama, University of Tokyo) was modified for TAP (pcpBpaRSv4). A Strep-tag, followed by a calmodulin-binding peptide (CBP), a V5 epitope, a tobacco etch virus (TEV) protease cleavage site, and a tandem protein A tag, was inserted after a mammalian expression promoter. Unique Rsr II and Sbf I restriction sites were placed between CBP and V5 epitope for subcloning SH3 domain coding sequences.

The sequences of SH3 domains inserted and the amino acid residues mutated to amber codon are as follows: ARHGAP26 (amino acids 759 to 814, UniProt Q9UNA1, Ser⁷⁷³), ARHGAP10 (amino acids 731 to 786, UniProt A1A4S6, Ser⁷⁴⁵), ARHGAP10L (amino acids 820 to 874, UniProt A6N128, His⁸³³), ARHGAP9 (amino acids 25 to 87, UniProt Q9BRR9, Asp⁶⁰), SNX26 (amino acids 189 to 247, UniProt O14559, Asp²⁰³), SRGAP2 (amino acids 725 to 786, UniProt O75044, Arg⁷⁴⁵), ARHGAP4 (amino acids 749 to 804, UniProt P98171, Gln⁷⁶³), RICS (amino acids 262 to 320, UniProt A7KAX9, Asp²⁷⁶), and ARHGAP12 (amino acids 15 to 73, UniProt Q8IWW6, Asp⁴⁹). N-terminal GST fusion of the SH3 domains was made by subcloning an SH3 domain into pGEX-4T vector or cloning into pDEST15 vector with GATEWAY system (Invitrogen). GFP-tagged gephyrin and palladin, Flag-tagged ARHGAP4 and CDC42BPA, and Myc-tagged SNX26 were provided by M. Kneussel (University of Hamburg, Germany), C. Otey (University of North Carolina), T. Pawson (Mount Sinai Hospital, Canada), T. Leung (Institute of Molecular and Cell Biology, Singapore), and A. Saltiel (University of Michigan), respectively. WRP-V5, WRP(Δ SH3)-V5, GFP-WRP, and GFP-WRP(Δ SH3) constructs were previously described (34). SRGAP2-Flag and SRGAP2-V5 were also described (33). Gephyrin-V5 and MICAL-V5 were made by subcloning the open reading frames (ORFs) of mRNA sequences (BC030016 and NM022765, respectively) into pcDNA3.1-V5/His (Invitrogen). To make GFP-tagged ARHGAP26 and WASF2, we subcloned the ORF sequences into pEGFP-C2 vector with ARHGAP26 mRNA (BC068555) and the WASF2 plasmid previously described (73), respectively. All the amber mutants were generated following the QuikChange site-directed mutagenesis protocol (Stratagene).

Photo-cross-linking

For photo-cross-linking of the SRGAP2 SH3 domain variants (wild-type, R745amber, or D764amber) in Chinese hamster ovary (CHO) cells, the cells were transfected using Lipofectamine 2000 (Invitrogen) grown for 24 hours in growth medium containing 1 mM pBpa and exposed to UV light for 30 min with an 8-W handheld UV light before cell lysis. For photo-cross-linking of Rho family GAP SH3 domains in FreeStyle 293 cells, the following method was used. For transfection, a 1-liter culture of FreeStyle 293 cells was grown in FreeStyle 293 medium (Gibco) to the density of 1×10^6 cells/ml. Transfection was done following a modified polyethylenimine (PEI) protocol (74). One milligram of the vector DNA encoding TAP-tagged Rho family GAP SH3 domain plus *Eco*-pBpaRS and suppressor tRNA^{Tyr} was mixed with 2 mg of PEI in 50 ml of FreeStyle 293 medium and added to the cell culture after 10 min of incubation. pBpa (270 mg) was dissolved in 1.1 ml of 1 M NaOH, filtered, and added to the cell culture dropwise after the addition of 7.5 ml of 1 M Hepes buffer

to the culture. After a 3-day incubation at 37°C in the dark, the cells were transferred into a homemade cross-linking chamber, which was a commercially available "UV pond clarifier" (Garden Treasures, Home Depot) whose standard 254-nm UV bulb was replaced with a black light bulb (peak wavelength, 365 nm). After a 30-min photo-cross-linking in the chamber, cells were pelleted by centrifugation and either stored at -80°C or used for TAP purification.

Tandem affinity purification

The cross-linked cell pellet was resuspended in 20 ml of TAP lysis buffer [50 mM tris (pH 7.5), 150 mM NaCl, 1 mM EDTA, 7.5% glycerol, 25 mM NaF, 0.5% NP-40, 1 mM Na₃VO₄, 1 mM dithiothreitol (DTT), 1 mM 4-(2-aminoethyl)benzenesulfonyl fluoride hydrochloride (AEBSE), leupeptin/pepstatin (2 μ g/ml)], homogenized with a Dounce homogenizer, and subjected to centrifugation at 17,200g for 30 min. The supernatant was further clarified by ultracentrifugation at 100,000g for 1 hour. The supernatant was mixed with prewashed immunoglobulin G (IgG)-Sephacrose beads (200- μ l bed volume) and incubated for 3 hours to overnight at 4°C. The beads were pelleted, resuspended with 0.5 ml of TAP lysis buffer, applied onto a chromatography column, and allowed to pack by gravity. After being washed with 50 ml of TAP lysis buffer and 5 ml of TEV cleavage buffer [10 mM tris (pH 7.5), 100 mM NaCl, 0.5 mM EDTA, 1% Triton X-100, 1 mM DTT, 1 mM AEBSE, leupeptin/pepstatin (2 μ g/ml)], the beads were transferred into an Eppendorf tube, resuspended with 0.2 ml of TEV cleavage buffer, and incubated with 100 U of TEV protease for 3 hours at room temperature and overnight at 4°C. The whole solution was applied onto a new chromatography column, and the flow-through was collected. Residual eluate was further collected by applying 150 μ l of TEV cleavage buffer four times to the column. Elution fractions were combined and mixed with prewashed calmodulin-Sepharose beads (400- μ l bed volume) in 1 ml of 2 \times CBP-binding buffer [10 mM tris (pH 7.5), 100 mM NaCl, 2 mM MgAc, 4 mM CaCl₂, 2 mM imidazole, 0.1% NP-40, 1 mM DTT, 1 mM AEBSE, leupeptin/pepstatin (2 μ g/ml)], and incubated for 2 hours to overnight at 4°C. The mixture was applied onto a new chromatography column. After being washed with 20 ml of calmodulin-rinsing buffer [50 mM ammonium bicarbonate (pH 8.0), 75 mM NaCl, 1 mM MgAc, 2 mM CaCl₂, 1 mM imidazole, 1 mM AEBSE, leupeptin/pepstatin (2 μ g/ml)] and 2 ml of 50 mM ammonium bicarbonate, the column was incubated with prewarmed calmodulin-elution buffer [50 mM ammonium bicarbonate (pH 8.0), 25 mM EGTA] containing 0.1% RapiGest for 10 min and eluted. This was repeated three times with calmodulin-elution buffer without RapiGest for 5 min. The eluates were combined and concentrated with an Amicon Ultra-4 [30,000 molecular weight cutoff (MWCO)] centrifugal filter device (Millipore) by centrifugation at 2500g for 5 min followed by the addition of 4 ml of 50 mM ammonium bicarbonate and centrifugation at 2500g for 10 min. The final sample contained the purified bait SH3 domain as well as the cross-linked species. One advantage of the relatively small SH3 domain, when compared to full-length bait proteins, is that the number of trypsin peptides in the sample from the bait domain was limited. This prevented the obscuring of the prey peptides by excess bait peptides during the subsequent MS identifications.

Sample preparation and nanoflow liquid chromatography electrospray ionization MS/MS analysis

After a total protein quantitation by a Bradford assay (Bio-Rad), samples were reduced with 5 mM DTT for 30 min at 70°C, and free sulfhydryls were alkylated with 10 mM iodoacetamide for 45 min at room temperature. Proteolytic digestion was accomplished by the addition of 500 ng of sequencing-grade trypsin (Promega) and incubation at 37°C for 18 hours. Samples were then acidified to pH 2.5 with trifluoroacetic acid (TFA) and incubated at 60°C

for 1 hour to hydrolyze remaining RapiGest surfactant. Insoluble hydrolyzed surfactant was cleared by centrifugation at 15,000 rpm for 5 min.

Chromatographic separation of peptide mixtures was performed on a Waters nanoACQUITY UPLC equipped with a 1.7- μ m BEH130 C₁₈ reversed-phase column [75 μ m inside diameter (ID) \times 250 mm]. The mobile phase consisted of (A) 0.1% formic acid in water and (B) 0.1% formic acid in acetonitrile. After a 5- μ l injection (1 μ g of total peptide), peptides were trapped for 5 min on a 5- μ m Symmetry C₁₈ column (180 μ m ID \times 20 mm) at 20 μ l/min in 99.9% A. The analytical column was then switched inline and a linear elution gradient of 5% B to 40% B was performed over 90 min at 300 nl/min. The analytical column was connected to a fused silica PicoTip emitter (New Objective) with a 10- μ m tip orifice and coupled to the mass spectrometer through an electrospray interface.

MS data were acquired on an LTQ-Orbitrap XL (TAP pull-downs and FreeStyle 293 lysate) (Thermo Scientific), QToF Premier (TAP pull-downs) (Waters), or QToF Ultima (TAP pull-downs) (Waters) mass spectrometer operating in positive-ion electrospray ionization mode. For data acquired on the LTQ-Orbitrap XL, the instrument was set to acquire a precursor MS scan in the Orbitrap from mass/charge ratio (m/z) 400 to 2000 with $r = 60,000$ at m/z 400 and a target AGC setting of 1×10^6 ions. In a data-dependent mode of acquisition, MS/MS spectra of either the five (TAP pull-down) or two (293 lysate) most abundant precursor ions were acquired in the Orbitrap at $r = 7500$ at m/z 400 with a target AGC setting of 2×10^5 ions. Maximum fill times were set to 1000 ms for full MS scans and 500 ms for MS/MS scans with minimum MS/MS triggering thresholds of 5000 counts. For all experiments, fragmentation occurred in the LTQ linear ion trap with a collision-induced dissociation (CID) energy setting of 35% and a dynamic exclusion of 60 s was used for previously fragmented precursor ions. For data acquired on the QToF Premier or QToF Ultima, the instrument was set to acquire a precursor MS scan from m/z 50 to 1990 followed by three data-dependent MS/MS product ion scans from m/z 50 to 1990 with a charge state-dependent CID energy setting. To increase coverage of lower-abundance precursor ions, we used a 120-s dynamic exclusion list. A separate LC channel with Glu-1-fibrinopeptide (200 fmol/ μ l) in 50% acetonitrile/0.1% formic acid flowing at 500 nl/min was referenced every 30 s through a nano lock-spray interface.

Qualitative identifications from raw LC-MS/MS data

Raw LC-MS/MS data files were processed in Mascot distiller (Matrix Science) and then submitted to independent Mascot database searches (Matrix Science) against SwissProt and National Center for Biotechnology Information (NCBI) databases (*Homo sapiens* taxonomy) containing both forward and reverse entries of each protein. Search tolerances for LTQ-Orbitrap XL data were between 5 and 10 parts per million (ppm) for precursor ions and 0.02 dalton for product ions depending on the instrument performance as determined by a system suitability test of 50 fmol of yeast alcohol dehydrogenase digest run immediately before each sample. Search tolerances for QToF Premier and QToF Ultima data were 20 ppm for precursor ions and 0.04 dalton for product ions. All data were searched using trypsin specificity with up to two missed cleavages. Carbamidomethylation (+57.0214 daltons on C) was set as a fixed modification, whereas oxidation (+15.9949 daltons on M) was considered a variable modification. All searched spectra were imported into Scaffold (Proteome Software), and low-stringency protein confidence thresholds were set with a Bayesian statistical algorithm based on the PeptideProphet and ProteinProphet algorithms, which yielded a peptide and protein FDR of 1.6% and 14.2%, respectively (75, 76).

Hierarchical clustering

Relative protein abundance in the sample prepared by TAP was quantified with spectral counting (16). We used “quantitative values” calculated with-

in Scaffold to provide a first-pass normalization of spectral counts for each protein based on the average total spectra counts across multiple samples (17). To further normalize the relative protein abundance, we expressed the (modified) spectral counts as a percentage of the total spectra observed in the sample. Mean normalized spectral counts were obtained from multiple independent experiments ($n \geq 2$ for each Rho family GAP). Hierarchical clustering was performed on the basis of the uncentered Pearson correlation of the mean normalized spectral counts by means of the Cluster 3.0 program (<http://bonsai.ims.utokyo.ac.jp/~mdehoon/software/cluster/>) (77). The dendrograms were viewed with the Java TreeView program (<http://jtreeview.sourceforge.net/>) (78).

Peptide array synthesis and SH3 domain overlay

Peptides (18-mer) were synthesized as previously described (34) with Auto Spot Robot ASP 222 (Intavis AG). GST-tagged SH3 domain overlays were performed with 100 μ M protein as previously described (73). Immunopositive peptide spots were densitometrically quantified with MetaMorph (version 7.7) (<http://www.moleculardevices.com/>).

Construction of interactomes

Rho family GAP protein-protein interaction networks were constructed by combining the SH3 domain–ligand interactions revealed through this study and previously known (physical) interactions. The known (physical) interactions were searched among (i) proteins identified in the PI clusters predominantly associated with individual Rho family GAP SH3 domains, (ii) subspecific proteins that showed association to two Rho family GAP SH3 domains only (including the focused Rho family GAP SH3 domain), and (iii) GTPases (RhoA, Cdc42, Rac1), using GeneMANIA (version 2.0) (<http://genemania.org/>), Ingenuity (<http://www.ingenuity.com/>), and STRING (version 8.3) (<http://string-db.org/>) (79). The open-source platform Cytoscape (version 2.8.0) was used to visualize the protein-protein networks (<http://www.cytoscape.org/>).

Immunostaining of cell lines and cultured neurons

HEK293 cells were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and penicillin/streptomycin. The cells were plated onto poly-D-lysine-coated glass coverslips and transfected with GFP-gephyrin and vector or WRP-V5 by means of Lipofectamine 2000 (Invitrogen), following the manufacturer's instruction. Two days after transfection, cells were washed with phosphate-buffered saline (PBS) and fixed for 10 min with 4% paraformaldehyde in PBS. Cells were then permeabilized for 3 min with 0.1% Triton X-100 in PBS. After being washed with PBS, the cells were incubated for 1 hour with blocking solution (10% normal goat serum in PBS) and incubated overnight at 4°C with monoclonal anti-V5 (Invitrogen, 1:200) in a blocking buffer containing 0.1% Triton X-100. After extensive washing, cells were incubated for 1 hour with Alexa-labeled secondary antibodies (Molecular Probes). Fluorescence signals were visualized with Zeiss 710 confocal laser scanning system (Zeiss). Confocal images (10 \times magnification) were randomly taken, and the cells observed in the images were analyzed with MetaMorph to quantify the area and fluorescence intensity of gephyrin clusters. Total 683 (Vector) or 326 (WRP-V5) of gephyrin clusters were measured from three independent experiments. To analyze gephyrin clustering in the neurons, we prepared hippocampal neuronal cultures from postnatal day 1 pups and transfected them with Lipofectamine 2000 (Invitrogen) as described previously (58). Neurons transfected at DIV9 (day 9 in vitro) with GFP, GFP-WRP, or WRP(ASH3) were fixed at DIV12 and processed for immunostaining as described above. The primary antibodies used are anti-gephyrin monoclonal antibody (mAb) (mAb7a, SYNaptic SYstems, 1:300) and polyclonal anti-GFP antibody (Invitrogen, 1:400) or polyclonal

rabbit anti-VIAAT (vesicular GABA transporter) (SYnaptic SYstems, 1:500). Neurons that were positive for GFP were randomly selected [GFP, $n = 42$ cells from seven mice; GFP-WRP, $n = 27$ cells from four mice; GFP-WRP (Δ SH3), $n = 14$ cells from three mice], and the confocal images of neuronal dendrites ($63\times$ magnification) were taken on a Zeiss 710 confocal microscope. The length of dendritic segments was measured with ImageJ software (National Institutes of Health), and the number of gephyrin puncta in the dendritic segments overlapping with GFP signal was counted manually.

Immunoprecipitation experiments

Coimmunoprecipitation experiments were performed as previously described in detail (33, 34). Briefly, lysates were precleared before incubation with antibody and either agarose-conjugated protein A or G overnight at 4°C with rocking. Antibodies and bound proteins from extracts were precipitated by centrifugation and extensively washed. Bound proteins were eluted in SDS sample buffer.

Statistical analysis

Statistical analysis, Student's t test (two-tailed), was done with 2007 Office Excel version 12.

Animals

Description of the WRP knockout animals has previously been published (58). These mice contain a LoxP flanked exon 3, allowing for conditional deletion upon Cre recombinase expression. Mice used in this study were from previous crosses into a cytomegalovirus (CMV)-based Cre transgenic line to create germline deletion of the WRP exon 3. All mice were housed in Duke University's Division of Laboratory Animal Resources facilities, and all procedures were approved by the Duke University Institutional Animal Care and Use Committee and were in accordance with National Institutes of Health guidelines.

Immunostaining of brain sections

Preparation of brain sections was as previously described (80) with some modifications. Briefly, WRP^{+/+} and WRP^{-/-} P40 mice ($n = 3$ for each genotype) were anesthetized and decapitated. The brains were rapidly excised and frozen in dry ice. Cryostat sections ($30\ \mu\text{m}$ thick) were mounted onto gelatin-coated slides and briefly fixed with pre-cold 2% paraformaldehyde/PBS solution for 10 min. The sections were subjected to antigen retrieval in citrate buffer [10 mM citric acid and 0.05% Tween 20 (pH 6.0)] before being incubated overnight with antibodies against mAb7a (SYnaptic SYstems, 1:200) and polyclonal rabbit anti-GABA_A receptor $\gamma 2$ subunit (SYnaptic SYstems, 1:300). For each mouse, four brain slices were processed for immunostaining. Quantification of gephyrin and GABA receptor puncta densities was performed with MetaMorph on two images per slice of the CA1 region of the hippocampus taken at a $63\times$ magnification. The area and fluorescence intensity of puncta were quantified with "puncta analyzer" plug-in (written by B. Wark) for ImageJ (81). Adjacent sections to stain with antibodies against mAb7a and VIAAT (1:500) were treated in the same way.

SUPPLEMENTARY MATERIALS

www.sciencesignaling.org/cgi/content/full/4/201/rs13/DC1

Materials and Methods

Fig. S1. Expression levels of Rho family GAP SH3 domains and their interactors.

Fig. S2. Identification of SH3 domain binding proteins by peptide array-based in vitro binding assay and coimmunoprecipitation.

Fig. S3. Rho family GAP PI cluster interactomes and colocalization data.

Fig. S4. Gephyrin's interaction with WRP and colocalization with VIAAT.

Fig. S5. Endogenous gephyrin is highly colocalized with inhibitory synapse presynaptic marker VIAAT in vivo.

Reference

REFERENCES AND NOTES

1. L. Van Aelst, C. D'Souza-Schoorey, Rho GTPases and signaling networks. *Genes Dev.* **11**, 2295–2322 (1997).
2. M. Symons, J. Settleman, Rho family GTPases: More than simple switches. *Trends Cell Biol.* **10**, 415–419 (2000).
3. J. Tcherkezian, N. Lamarche-Vane, Current knowledge of the large RhoGAP family of proteins. *Biol. Cell* **99**, 67–86 (2007).
4. A. Bernards, J. Settleman, GAP control: Regulating the regulators of small GTPases. *Trends Cell Biol.* **14**, 377–385 (2004).
5. B. J. Mayer, SH3 domains: Complexity in moderation. *J. Cell Sci.* **114**, 1253–1263 (2001).
6. S. S. Li, Specificity and versatility of SH3 and other proline-recognition domains: Structural basis and implications for cellular signal transduction. *Biochem. J.* **390**, 641–653 (2005).
7. J. M. Robens, L. Yeow-Fong, E. Ng, C. Hall, E. Manser, Regulation of IRSp53-dependent filopodial dynamics by antagonism between 14-3-3 binding and SH3-mediated localization. *Mol. Cell. Biol.* **30**, 829–844 (2010).
8. J. C. Kauer, S. Erickson-Viitanen, H. R. Wolfe Jr., W. F. DeGrado, *p*-Benzoyl-L-phenylalanine, a new photoreactive amino acid. Photolabeling of calmodulin with a synthetic calmodulin-binding peptide. *J. Biol. Chem.* **261**, 10695–10700 (1986).
9. K. T. O'Neil, S. Erickson-Viitanen, W. F. DeGrado, Photolabeling of calmodulin with basic, amphiphilic α -helical peptides containing *p*-benzoylphenylalanine. *J. Biol. Chem.* **264**, 14571–14578 (1989).
10. N. Hino, Y. Okazaki, T. Kobayashi, A. Hayashi, K. Sakamoto, S. Yokoyama, Protein photo-cross-linking in mammalian cells by site-specific incorporation of a photoreactive amino acid. *Nat. Methods* **2**, 201–206 (2005).
11. I. S. Farrell, R. Toroney, J. L. Hazen, R. A. Mehl, J. W. Chin, Photo-cross-linking interacting proteins with a genetically encoded benzophenone. *Nat. Methods* **2**, 377–384 (2005).
12. W. Wang, J. K. Takimoto, G. V. Louie, T. J. Baiga, J. P. Noel, K. F. Lee, P. A. Slesinger, L. Wang, Genetically encoding unnatural amino acids for cellular and neuronal studies. *Nat. Neurosci.* **10**, 1063–1072 (2007).
13. W. Liu, A. Brock, S. Chen, P. G. Schultz, Genetic incorporation of unnatural amino acids into proteins in mammalian cells. *Nat. Methods* **4**, 239–244 (2007).
14. K. Sakamoto, A. Hayashi, A. Sakamoto, D. Kiga, H. Nakayama, A. Soma, T. Kobayashi, M. Kitabatake, K. Takio, K. Saito, M. Shirouzu, I. Hirao, S. Yokoyama, Site-specific incorporation of an unnatural amino acid into proteins in mammalian cells. *Nucleic Acids Res.* **30**, 4692–4699 (2002).
15. X. Li, Y. Chen, Y. Liu, J. Gao, F. Gao, M. Bartlam, J. Y. Wu, Z. Rao, Structural basis of Robo proline-rich motif recognition by the srGAP1 Src homology 3 domain in the Slit-Robo signaling pathway. *J. Biol. Chem.* **281**, 28430–28437 (2006).
16. H. Liu, R. G. Sadygov, J. R. Yates III, A model for random sampling and estimation of relative protein abundance in shotgun proteomics. *Anal. Chem.* **76**, 4193–4201 (2004).
17. B. C. Searle, Scaffold: A bioinformatic tool for validating MS/MS-based proteomic studies. *Proteomics* **10**, 1265–1269 (2010).
18. M. B. Eisen, P. T. Spellman, P. O. Brown, D. Botstein, Cluster analysis and display of genome-wide expression patterns. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 14863–14868 (1998).
19. R. Tonikian, X. Xin, C. P. Toret, D. Gfeller, C. Landgraf, S. Panni, S. Paoluzi, L. Castagnoli, B. Currell, S. Seshagiri, H. Yu, B. Winsor, M. Vidal, M. B. Gerstein, G. D. Bader, R. Volkmer, G. Cesareni, D. G. Drubin, P. M. Kim, S. S. Sidhu, C. Boone, Bayesian modeling of the yeast SH3 domain interactome predicts spatiotemporal dynamics of endocytosis proteins. *PLoS Biol.* **7**, e1000218 (2009).
20. J. D. Hildebrand, J. M. Taylor, J. T. Parsons, An SH3 domain-containing GTPase-activating protein for Rho and Cdc42 associates with focal adhesion kinase. *Mol. Cell. Biol.* **16**, 3169–3178 (1996).
21. H. Shibata, K. Oishi, A. Yamagiwa, M. Matsumoto, H. Mukai, Y. Ono, PKN β interacts with the SH3 domains of Grf and a novel Grf related protein, Grf2, which are GTPase activating proteins for Rho family. *J. Biochem.* **130**, 23–31 (2001).
22. J. F. Rual, K. Venkatesan, T. Hao, T. Hirozane-Kishikawa, A. Dricot, N. Li, G. F. Berriz, F. D. Gibbons, M. Dreze, N. Ayivi-Guedehoussou, N. Klitgord, C. Simon, M. Boxem, S. Milstein, J. Rosenberg, D. S. Goldberg, L. V. Zhang, S. L. Wong, G. Franklin, S. Li, J. S. Albala, J. Lim, C. Fraughton, E. Llamas, S. Cevik, C. Bex, P. Lamesch, R. S. Sikorski, J. Vandenhaute, H. Y. Zoghbi, A. Smolyar, S. Bosak, R. Sequerra, L. Doucette-Stamm, M. E. Cusick, D. E. Hill, F. P. Roth, M. Vidal, Towards a proteome-scale map of the human protein-protein interaction network. *Nature* **437**, 1173–1178 (2005).
23. X. R. Ren, Q. S. Du, Y. Z. Huang, S. Z. Ao, L. Mei, W. C. Xiong, Regulation of CDC42 GTPase by proline-rich tyrosine kinase 2 interacting with PSGAP, a novel pleckstrin homology and Src homology 3 domain containing rhoGAP protein. *J. Cell Biol.* **152**, 971–984 (2001).
24. A. Kondo, S. Hashimoto, H. Yano, K. Nagayama, Y. Mazaki, H. Sabe, A new paxillin-binding protein, PAG3/Papa/KIAA0400, bearing an ADP-ribosylation factor

- GTPase-activating protein activity, is involved in paxillin recruitment to focal adhesions and cell migration. *Mol. Biol. Cell* **11**, 1315–1327 (2000).
25. R. J. Hung, U. Yazdani, J. Yoon, H. Wu, T. Yang, N. Gupta, Z. Huang, W. J. van Berkel, J. R. Terman, Mical links semaphorins to F-actin disassembly. *Nature* **463**, 823–827 (2010).
 26. T. Higashi, T. Ikeda, T. Murakami, R. Shirakawa, M. Kawato, K. Okawa, M. Furuse, T. Kimura, T. Kita, H. Horiuchi, Flightless-I (Fli-I) regulates the actin assembly activity of diaphanous-related formins (DRFs) Daam1 and mDia1 in cooperation with active Rho GTPase. *J. Biol. Chem.* **285**, 16231–16238 (2010).
 27. E. B. Ott, N. M. van den Akker, P. A. Sakalis, A. C. Gittenberger-de Groot, A. J. Te Velthuis, C. P. Bagowski, The lim domain only protein 7 is important in zebrafish heart development. *Dev. Dyn.* **237**, 3940–3952 (2008).
 28. M. Matsuda, Y. Kobayashi, S. Masuda, M. Adachi, T. Watanabe, J. K. Yamashita, E. Nishi, S. Tsukita, M. Furuse, Identification of adherens junction-associated GTPase activating proteins by the fluorescence localization-based expression cloning. *Exp. Cell Res.* **314**, 939–949 (2008).
 29. G. Gamba, Role of WNK kinases in regulating tubular salt and potassium transport and in the development of hypertension. *Am. J. Physiol. Renal Physiol.* **288**, F245–F252 (2005).
 30. C. Tribioli, S. Droetto, S. Bione, G. Cesareni, M. R. Torrisi, L. V. Lotti, L. Lanfranccone, D. Toniolo, P. Pelicci, An X chromosome-linked gene encoding a protein with characteristics of a rhoGAP predominantly expressed in hematopoietic cells. *Proc. Natl. Acad. Sci. U.S.A.* **93**, 695–699 (1996).
 31. K. G. Campellone, M. D. Welch, A nucleator arms race: Cellular control of actin assembly. *Nat. Rev. Mol. Cell Biol.* **11**, 237–251 (2010).
 32. T. Takenawa, S. Suetsugu, The WASP–WAVE protein network: Connecting the membrane to the cytoskeleton. *Nat. Rev. Mol. Cell Biol.* **8**, 37–48 (2007).
 33. F. M. Mason, E. G. Heimsath, H. N. Higgs, S. H. Soderling, Bi-modal regulation of a formin by srGAP2. *J. Biol. Chem.* **286**, 6577–6586 (2011).
 34. S. H. Soderling, K. L. Binns, G. A. Wayman, S. M. Davee, S. H. Ong, T. Pawson, J. D. Scott, The WRP component of the WAVE-1 complex attenuates Rac-mediated signalling. *Nat. Cell Biol.* **4**, 970–975 (2002).
 35. Y. Furukawa, T. Kawasoe, Y. Daigo, T. Nishiwaki, H. Ishiguro, M. Takahashi, J. Kitayama, Y. Nakamura, Isolation of a novel human gene, *ARHGAP9*, encoding a Rho-GTPase activating protein. *Biochem. Biophys. Res. Commun.* **284**, 643–649 (2001).
 36. B. K. Ang, C. Y. Lim, S. S. Koh, N. Sivakumar, S. Taib, K. B. Lim, S. Ahmed, G. Rajagopal, S. H. Ong, ArhGAP9, a novel MAP kinase docking protein, inhibits Erk and p38 activation through WW domain binding. *J. Mol. Signal.* **2**, 1 (2007).
 37. G. Scita, P. Tenca, L. B. Arecas, A. Tocchetti, E. Frittoli, G. Giardina, I. Ponzanelli, P. Sini, M. Innocenti, P. P. Di Fiore, An effector region in Eps8 is responsible for the activation of the Rac-specific GEF activity of Sos-1 and for the proper localization of the Rac-based actin-polymerizing machine. *J. Cell Biol.* **154**, 1031–1044 (2001).
 38. A. Disanza, S. Mantoani, M. Hertzog, S. Gerboth, E. Frittoli, A. Steffen, K. Berhoerster, H. J. Krienkamp, F. Milanese, P. P. Di Fiore, A. Ciliberto, T. E. Stradal, G. Scita, Regulation of cell shape by Cdc42 is mediated by the synergic actin-bundling activity of the Eps8–IRSp53 complex. *Nat. Cell Biol.* **8**, 1337–1347 (2006).
 39. S. H. Chiang, J. Hwang, M. Legendre, M. Zhang, A. Kimura, A. R. Saltiel, TCGAP, a multidomain Rho GTPase-activating protein involved in insulin-stimulated glucose transport. *EMBO J.* **22**, 2679–2691 (2003).
 40. M. Rosário, R. Franke, C. Bednarski, W. Birchmeier, The neurite outgrowth multi-adaptor RhoGAP, NOMA-GAP, regulates neurite extension through SHP2 and Cdc42. *J. Cell Biol.* **178**, 503–516 (2007).
 41. H. Liu, T. Nakazawa, T. Tezuka, T. Yamamoto, Physical and functional interaction of Fyn tyrosine kinase with a brain-enriched Rho GTPase-activating protein TCGAP. *J. Biol. Chem.* **281**, 23611–23619 (2006).
 42. H. Yano, F. Cong, R. B. Birge, S. P. Goff, M. V. Chao, Association of the Abl tyrosine kinase with the Trk nerve growth factor receptor. *J. Neurosci. Res.* **59**, 356–364 (2000).
 43. M. Escalante, J. Courtney, W. G. Chin, K. K. Teng, J. I. Kim, J. E. Fajardo, B. J. Mayer, B. L. Hempstead, R. B. Birge, Phosphorylation of c-Crk II on the negative regulatory Tyr²²² mediates nerve growth factor-induced cell spreading and morphogenesis. *J. Biol. Chem.* **275**, 24787–24797 (2000).
 44. S. Wilkinson, H. F. Paterson, C. J. Marshall, Cdc42-MRCK and Rho-ROCK signalling cooperate in myosin phosphorylation and cell invasion. *Nat. Cell Biol.* **7**, 255–261 (2005).
 45. K. Moriyama, I. Yahara, Human CAP1 is a key factor in the recycling of cofilin and actin for rapid actin turnover. *J. Cell Sci.* **115**, 1591–1601 (2002).
 46. J. Papin, S. Subramaniam, Bioinformatics and cellular signaling. *Curr. Opin. Biotechnol.* **15**, 78–81 (2004).
 47. M. Nozumi, T. Togano, K. Takahashi-Niki, J. Lu, A. Honda, M. Taoka, T. Shinkawa, H. Koga, K. Takeuchi, T. Isobe, M. Igarashi, Identification of functional marker proteins in the mammalian growth cone. *Proc. Natl. Acad. Sci. U.S.A.* **106**, 17211–17216 (2009).
 48. J. Lu, M. Nozumi, K. Takeuchi, H. Abe, M. Igarashi, Expression and function of neuronal growth-associated proteins (nGAPs) in PC12 cells. *Neurosci. Res.* **70**, 85–90 (2011).
 49. Z. Wills, M. Emerson, J. Rusch, J. Bikoff, B. Baum, N. Perrimon, D. Van Vactor, A *Drosophila* homolog of cyclase-associated proteins collaborates with the Abl tyrosine kinase to control midline axon pathfinding. *Neuron* **36**, 611–622 (2002).
 50. T. Nakazawa, A. M. Watabe, T. Tezuka, Y. Yoshida, K. Yokoyama, H. Umemori, A. Inoue, S. Okabe, T. Manabe, T. Yamamoto, p250GAP, a novel brain-enriched GTPase-activating protein for Rho family GTPases, is involved in the *N*-methyl-D-aspartate receptor signaling. *Mol. Biol. Cell* **14**, 2921–2934 (2003).
 51. S. Impey, M. Davare, A. Lasiek, D. Fortin, H. Ando, O. Varlamova, K. Obrietan, T. R. Soderling, R. H. Goodman, G. A. Wayman, An activity-induced microRNA controls dendritic spine formation by regulating Rac1-PAK signaling. *Mol. Cell. Neurosci.* **43**, 146–156 (2010).
 52. G. A. Wayman, M. Davare, H. Ando, D. Fortin, O. Varlamova, H. Y. Cheng, D. Marks, K. Obrietan, T. R. Soderling, R. H. Goodman, S. Impey, An activity-regulated microRNA controls dendritic plasticity by down-regulating p250GAP. *Proc. Natl. Acad. Sci. U.S.A.* **105**, 9093–9098 (2008).
 53. R. Dhavan, L. H. Tsai, A decade of CDK5. *Nat. Rev. Mol. Cell Biol.* **2**, 749–759 (2001).
 54. Q. Huang, A. Raya, P. DeJesus, S. H. Chao, K. C. Quon, J. S. Caldwell, S. K. Chanda, J. C. Izpisua-Belmonte, P. G. Schultz, Identification of p53 regulators by genome-wide functional analysis. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 3456–3461 (2004).
 55. J. M. Fritschy, R. J. Harvey, G. Schwarz, Gephyrin: Where do we stand, where do we go? *Trends Neurosci.* **31**, 257–264 (2008).
 56. J. Kirsch, H. Betz, The postsynaptic localization of the glycine receptor-associated protein gephyrin is regulated by the cytoskeleton. *J. Neurosci.* **15**, 4148–4156 (1995).
 57. M. Bausen, J. C. Fuhrmann, H. Betz, G. A. O'Sullivan, The state of the actin cytoskeleton determines its association with gephyrin: Role of ena/VASP family members. *Mol. Cell. Neurosci.* **31**, 376–386 (2006).
 58. B. R. Carlson, K. E. Lloyd, A. Kruszewski, I. H. Kim, R. M. Rodriguez, C. Heindel, M. Faytell, S. M. Dudek, W. C. Wetsel, S. H. Soderling, WRP/srGAP3 facilitates the initiation of spine development by an inverse F-BAR domain, and its loss impairs long-term memory. *J. Neurosci.* **31**, 2447–2460 (2011).
 59. E. J. Marco, F. E. Abidi, J. Bristow, W. B. Dean, P. Cotter, R. J. Jeremy, C. E. Schwartz, E. H. Sherr, ARHGEF9 disruption in a female patient is associated with X linked mental retardation and sensory hyperarousal. *J. Med. Genet.* **45**, 100–105 (2008).
 60. V. M. Kalscheuer, L. Musante, C. Fang, K. Hoffmann, C. Fuchs, E. Carta, E. Deas, K. Venkateswarlu, C. Menzel, R. Ullmann, N. Tommerup, L. Dalprà, A. Tzschach, A. Selicorni, B. Lüscher, H. H. Ropers, K. Harvey, R. J. Harvey, A balanced chromosomal translocation disrupting ARHGEF9 is associated with epilepsy, anxiety, aggression, and mental retardation. *Hum. Mutat.* **30**, 61–68 (2009).
 61. S. Kins, H. Betz, J. Kirsch, Collybistin, a newly identified brain-specific GEF, induces submembrane clustering of gephyrin. *Nat. Neurosci.* **3**, 22–29 (2000).
 62. A. Pouloupoulos, G. Aramuni, G. Meyer, T. Soykan, M. Hoon, T. Papadopoulos, M. Zhang, I. Paarmann, C. Fuchs, K. Harvey, P. Jedlicka, S. W. Schwarzacher, H. Betz, R. J. Harvey, N. Brose, W. Zhang, F. Varoqueaux, Neurologin 2 drives postsynaptic assembly at perisomatic inhibitory synapses through gephyrin and collybistin. *Neuron* **63**, 628–642 (2009).
 63. T. Papadopoulos, M. Korte, V. Eulenburg, H. Kubota, M. Retiounskaia, R. J. Harvey, K. Harvey, G. A. O'Sullivan, B. Laube, S. Hulsman, J. R. Geiger, H. Betz, Impaired GABAergic transmission and altered hippocampal synaptic plasticity in collybistin-deficient mice. *EMBO J.* **26**, 3888–3899 (2007).
 64. H. S. Lee, R. D. Dimla, P. G. Schultz, Protein-DNA photo-crosslinking with a genetically encoded benzophenone-containing amino acid. *Bioorg. Med. Chem. Lett.* **19**, 5222–5224 (2009).
 65. A. C. Gingras, R. Aebersold, B. Raught, Advances in protein complex analysis using mass spectrometry. *J. Physiol.* **563**, 11–21 (2005).
 66. L. Armstrong, M. Lako, I. van Herpe, J. Evans, G. Saretzki, N. Hole, A role for nucleoprotein Zap3 in the reduction of telomerase activity during embryonic stem cell differentiation. *Mech. Dev.* **121**, 1509–1522 (2004).
 67. K. Harvey, I. C. Duguid, M. J. Aldred, S. E. Beatty, H. Ward, N. H. Keep, S. E. Lingenfelter, B. R. Pearce, J. Lundgren, M. J. Owen, T. G. Smart, B. Luscher, M. I. Rees, R. J. Harvey, The GDP-GTP exchange factor collybistin: An essential determinant of neuronal gephyrin clustering. *J. Neurosci.* **24**, 5816–5826 (2004).
 68. S. Reddy-Alla, B. Schmitt, J. Birkenfeld, V. Eulenburg, S. Dutertre, C. Böhringer, M. Götz, H. Betz, T. Papadopoulos, PH-domain-driven targeting of collybistin but not Cdc42 activation is required for synaptic gephyrin clustering. *Eur. J. Neurosci.* **31**, 1173–1184 (2010).
 69. V. Endris, B. Wogatzky, U. Leimer, D. Bartsch, M. Zatyka, F. Latif, E. R. Maher, G. Tariverdian, S. Kirsch, D. Karch, G. A. Rappold, The novel Rho-GTPase activating gene MEGAP/srGAP3 has a putative role in severe mental retardation. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 11754–11759 (2002).

70. A. M. Addington, J. L. Rapoport, The genetics of childhood-onset schizophrenia: When madness strikes the prepubescent. *Curr. Psychiatry Rep.* **11**, 156–161 (2009).
71. D. A. Lewis, G. González-Burgos, Neuroplasticity of neocortical circuits in schizophrenia. *Neuropsychopharmacology* **33**, 141–165 (2008).
72. T. Enomoto, M. T. Tse, S. B. Floresco, Reducing prefrontal γ -aminobutyric acid activity induces cognitive, behavioral, and dopaminergic abnormalities that resemble schizophrenia. *Biol. Psychiatry* **69**, 432–441 (2011).
73. R. S. Westphal, S. H. Soderling, N. M. Alto, L. K. Langeberg, J. D. Scott, Scar/WAVE-1, a Wiskott-Aldrich syndrome protein, assembles an actin-associated multi-kinase scaffold. *EMBO J.* **19**, 4589–4600 (2000).
74. Y. Durocher, S. Perret, A. Kamen, High-level and high-throughput recombinant protein production by transient transfection of suspension-growing human 293-EBNA1 cells. *Nucleic Acids Res.* **30**, E9 (2002).
75. A. Keller, A. I. Nesvizhskii, E. Kolker, R. Aebersold, Empirical statistical model to estimate the accuracy of peptide identifications made by MS/MS and database search. *Anal. Chem.* **74**, 5383–5392 (2002).
76. A. I. Nesvizhskii, A. Keller, E. Kolker, R. Aebersold, A statistical model for identifying proteins by tandem mass spectrometry. *Anal. Chem.* **75**, 4646–4658 (2003).
77. M. J. de Hoon, S. Imoto, J. Nolan, S. Miyano, Open source clustering software. *Bioinformatics* **20**, 1453–1454 (2004).
78. A. J. Saldanha, Java Treeview—Extensible visualization of microarray data. *Bioinformatics* **20**, 3246–3248 (2004).
79. D. Szklarczyk, A. Franceschini, M. Kuhn, M. Simonovic, A. Roth, P. Minguéz, T. Doerks, M. Stark, J. Muller, P. Bork, L. J. Jensen, C. von Mering, The STRING database in 2011: Functional interaction networks of proteins, globally integrated and scored. *Nucleic Acids Res.* **39**, D561–D568 (2011).
80. E. M. Schneider Gasser, C. J. Straub, P. Panzanelli, O. Weinmann, M. Sassoè-Pognetto, J. M. Fritschy, Immunofluorescence in brain sections: Simultaneous detection of pre-synaptic and postsynaptic proteins in identified neurons. *Nat. Protoc.* **1**, 1887–1897 (2006).
81. D. M. Ippolito, C. Eroglu, Quantifying synapses: An immunocytochemistry-based assay to quantify synapse number. *J. Vis. Exp.* 10.3791/2270 (2010).

Acknowledgments: We thank M. Sassoè-Pognetto (University of Turin) for the advice on immunohistochemistry; S. Yokoyama (University of Tokyo), M. Kneussel (University of Hamburg), C. Otey (University of North Carolina), T. Pawson (Mount Sinai Hospital), T. Leung (Institute of Molecular and Cell Biology), and A. Saltiel (University of Michigan) for the gifts of plasmids. We also thank W. Thompson and M. Turner (Duke University) for help with the lysate expression analysis and for MS sample preparation. **Funding:** This work was supported by NIH grants R21-CA-140030–01 and R01-NS059957 (S.H.S.). **Author contributions:** S.H.S. and H.O. conceived the project; H.O., F.M.M., E.J.S., and A.U. performed the experiments and analyzed the data; S.H.S. supervised the entire project; H.O. and S.H.S. wrote the manuscript; E.J.S., M.A.M., and F.M.M. commented on and edited the manuscript. **Competing interests:** The authors declare that they have no competing financial interests.

Submitted 13 May 2011

Accepted 4 November 2011

Final Publication 29 November 2011

10.1126/scisignal.2002189

Citation: H. Okada, A. Uezu, F. M. Mason, E. J. Soderblom, M. A. Moseley III, S. H. Soderling, SH3 domain-based phototrapping in living cells reveals Rho family GAP signaling complexes. *Sci. Signal.* **4**, rs13 (2011).

Supplementary Materials for

SH3 Domain–Based Phototrapping in Living Cells Reveals Rho Family GAP Signaling Complexes

Hirokazu Okada, Akiyoshi Uezu, Frank M. Mason, Erik J. Soderblom, M. Arthur Moseley III, Scott H. Soderling*

*To whom correspondence should be addressed. E-mail: s.soderling@cellbio.duke.edu

Published 29 November 2011, *Sci. Signal.* **4**, rs13 (2011)
DOI: 10.1126/scisignal.2002189

The PDF file includes:

Materials and Methods

Fig. S1. Expression levels of Rho family GAP SH3 domains and their interactors.

Fig. S2. Identification of SH3 domain binding proteins by peptide array–based in vitro binding assay and coimmunoprecipitation.

Fig. S3. Rho family GAP PI cluster interactomes and colocalization data.

Fig. S4. Gephyrin's interaction with WRP and colocalization with VIAAT.

Fig. S5. Endogenous gephyrin is highly colocalized with inhibitory synapse presynaptic marker VIAAT in vivo.

Reference

Supplementary Materials

SUPPLEMENTARY MATERIALS AND METHODS

Label-Free Quantitation of Free Style 293 lysate

Label-free quantitation and integration of qualitative peptide identifications was performed using Rosetta Elucidator (v 3.3, Rosetta Inpharmatics, Seattle, WA). Triplicate LC-MS/MS analysis of a Free Style 293 lysate were imported and subjected to chromatographic retention time alignment using the PeakTeller® algorithm. Quantitation of all detectable signals in the precursor MS spectra was performed within Elucidator by calculating peak height of the corresponding peptide level extracted ion chromatograms. Protein quantities were calculated using the average MS response from the two or three highest abundant peptides based on a modified strategy initially described by Silva et. al. (1). Absolute quantities were determined by normalizing MS response factors to that of yeast alcohol dehydrogenase spiked into the lysate at 50 fmol/ug.

Immunostaining of HBE cells

Culturing and transfection of Human Bronchial Epithelial (HBE) cells were as described for HEK293 cells. Immunostaining for HBE cells was also performed the same way as for HEK293 cells except for the use of 1% paraformaldehyde for fixation. The primary antibodies used for staining were polyclonal mouse anti-ARHGAP12 (Abcam, 1:100) and rabbit anti-ZO-2 (C-term) (Invitrogen, 1:100).

REFERENCE

1. J. C. Silva, M. V. Gorenstein, G. Z. Li, J. P. Vissers, S. J. Geromanos, Absolute quantification of proteins by LCMSE: a virtue of parallel MS acquisition. *Mol Cell Proteomics* **5**, 144-156 (2006).

Fig. S1. Expression levels of Rho family GAP SH3 domains and their interactors. **(A)** Schematic of vector used to express pBpa containing bait SH3 domains. **(B)** Relative levels of each SH3 domain bait after transfection. n=2. **(C)** Graph depicting the quantification of the most abundant proteins in the lysate of Free Style 293 cells to three orders of magnitude. **(D)** Chart showing the number of PI cluster proteins identified in lysates. Only 10.8% of PI cluster proteins and one SH3 domain ligand were identified as abundant proteins in (C) using label-free LC/MS/MS quantification of Free Style 293 lysates.

Figure S1





Fig. S2. Identification of SH3 domain binding proteins by peptide array-based in vitro binding assay and coimmunoprecipitation. **A-I)** Proteins in Rho GAP SH3 domain-specific protein interaction (PI) clusters shown in Fig. 3A were searched for PXXP motifs. Peptides (18mer) containing the PXXP motifs were synthesized on cellulose membranes, incubated with purified GST-tagged wild-type Rho GAP SH3 domain, and immunoblotted with anti-GST antibody. The peptide array immunoblots were densitometrically quantified. The binding strength was normalized to the strongest interaction. **J)** HEK293 cells co-expressing palladin-GFP with vector or full-length wild-type SRGAP2-Flag were subjected to immunoprecipitation by Flag antibody. **K)** HEK293 cells co-expressing ARHGAP4 -Flag with GFP only or GFP-tagged full-length DIAPH1, were subjected to immunoprecipitation by GFP antibody. Co-precipitation was analyzed by Western blot analyses using indicated antibodies. Representative blots are shown from n=3.

B

Figure S2. ARHGAP10/SH3 - *in Vitro* Binding Assay



C


Figure S2. ARHGAP10L/SH3 - *in Vitro* Binding Assay

Figure S2. ARHGAP9/SH3 - *in Vitro* Binding Assay



Figure S2. SNX26/SH3 - *in Vitro* Binding Assay


E



Figure S2. SRGAP2/SH3 - *in Vitro* Binding Assay (1/2)



Figure S2. SRGAP2/SH3 - *in Vitro* Binding Assay (2/2)



0 20 40 60 80 100




Figure S2. ARHGAP4/SH3 - *in Vitro* Binding Assay (2/2)

G2




Figure S2. RICS/SH3 - *in Vitro* Binding Assay



Figure S2. ARHGAP12/SH3 - *in Vitro* Binding Assay



Figure S2. Co-immunoprecipitation assay





Fig. S3. Rho family GAP PI cluster interactomes and colocalization data. **A-I)** Protein-protein interaction networks of Rho GAPs were constructed from SH3 domain PI clusters based on peptide array and published interactome data. Circle nodes represent proteins in the PI clusters specific to individual RhoGAP SH3 domains. Hexagon nodes represent sub-specific proteins that were association with two RhoGAP SH3 domains. These two types of nodes are colored to reflect the mean normalized spectral counts (see spectral scale). Diamond nodes in cyan represent Rho GTPases added to visualize their potential links to the networks. Brown edges represent novel SH3-ligand interactions, and orange edges represent known interactions reproduced by this study. Thickness of these edges reflects the relative binding affinity determined by the in vitro binding assays. Blue and cyan edges that have a constant thickness indicate previously known interactions. Cyan edges exhibit known interactions with Rho GTPases. **J)** Co-localization of Endogenous ARHGAP12 and Tight Junction Protein TJP2 at cell-cell junctions. Human bronchial epithelial cells were fixed and stained with antibodies against indicated proteins. Scale bars = 10µm.

Figure S3. Rho GAP Interactome – ARHGAP26

A





Figure S3. Rho GAP Interactome – ARHGAP10

B



C




Figure S3. Rho GAP Interactome – ARHGAP9

D




Figure S3. Rho GAP Interactome – SNX26

E




Figure S3. Rho GAP Interactome – SRGAP2

F




Figure S3. Rho GAP Interactome - ARHGAP4

G




Figure S3. Rho GAP Interactome – RICS

H




Figure S3. Rho GAP Interactome – ARHGAP12

I




Figure S3. Co-localization Data

J




Fig. S4. Gephyrin's Interaction with WRP and colocalization with VIAAT. **A)** HEK293 cells co-expressing GFP-gephyrin with vector, V5-tagged full-length WRP or WRP lacking its SH3 domain, were subjected to immunoprecipitation by V5 antibody. Co-precipitation was analyzed by Western blot analysis using indicated antibodies. Representative blot from n=3 is shown. **B)** Co-localization of endogenous gephyrin and inhibitory synapse presynaptic marker VIAAT in hippocampal neurons. Dissociated hippocampal neurons were fixed at DIV12 and stained with gephyrin antibody mAb7a and VIAAT antibody. Scale bars = 10 μ m.




Fig. S5. Endogenous gephyrin is highly colocalized with inhibitory synapse presynaptic marker VIAAT in vivo. **A)** Stratum radiatum of CA1 region in the hippocampal formation from P40 WRP^{+/+} and WRP^{-/-} mice was stained with mAb7a and VIAAT antibodies. Scale bars = 10µm. **B)** Quantification of the gephyrin and VIAAT puncta density. n=3 mice. From each mouse, 4 brain slices were processed for immunostaining and 2 images/slice were obtained. The data are represented as mean ± SEM.

